

Child Health and the Income Gradient in Australia: The Role of Chronic Conditions

Rasheda Khanam^{a*}, Hong Son Nghiem^b and Luke B. Connelly^{b,c,d}

^aSchool of Accounting, Economics and Finance, University of Southern Queensland

^bCentre of National Research on Disability and Rehabilitation Medicine (CONROD)

^cAustralian Centre for Economic Research on Health

^dSchool of Economics, The University of Queensland

Abstract

In this paper we utilise data from the Longitudinal Study of Australian (LSAC) to examine the question of how the income-child health gradient, which has attracted much attention from the researchers in recent years, may be explained. Our results show that a previous health shock, as measured by chronic conditions, increases the probability of a child being reported as being in poor health in current period. We find that income has a protective effect on child health, but our interaction term of family income and the lagged health shock is not statistically significant. Therefore, our results do not provide any evidence that low-SES households are less able to manage a health shock than the high-SES household. Interestingly, we also do not find any convincing evidence that low-SES Australian children are subject to more health shocks. We speculate that the extensive and universal public healthcare system in Australia may offset some of the important health-related disadvantages that are associated with low incomes in other countries.

Key words: Child health, Income gradient, Chronic condition, Panel data, Australia

JEL Classification: I1

*Corresponding author, khanam@usq.edu.au; Phone: 61 7 4631 1256; Fax: + 61 7 4631 5594.

1 Introduction

Recently the relationship between child health and household income has received considerable amount of interest from the researchers and policy makers. This interest was generated by an influential article by Case, Lubotsky, and Paxson (2002), and since then a large numbers of papers have been published from internationally reputed journals. Most of this literature are based on the USA data and a few of them are from other countries (for example, UK studies- Currie, Shields, and Price, 2007; Propper, Rigg, and Burgess, 2007; Canadian study - Currie and Stabile, 2003; Australian study - Khanam, Nghiem, and Connelly, 2009). This growing literature (Case, Lubotsky, and Paxson, 2002; Currie and Stabile, 2003; Currie, Shields, and Price, 2007, Khanam, Nghiem and Connelly, 2008; Condliffe and Link, 2008 and Murasko, 2008) has established that there is a positive relationship between child health and household income. Some studies, for example, Case et al (2002), Currie and Stabile (2003, subsequently C&S), Condliffe and Link (2008) and Murasko (2008) found that this relationship is more pronounced for older children. Khanam, Nghiem, Connelly (2008) provided the first Australian evidence on this issue using Longitudinal Study of Australian Children (LSAC). They found similar trend for Australian children (from early to mid child-hood) when similar variables as previous authors are used.

Case et al. (2002) explored the possible mechanisms of the positive relationship between income-child health and the increasing gradient in their cross-sectional American sample¹ and found no precise mechanism that could explain this relationship and the increasing income gradient. Currie and Stabile (2003)'s study was the first that provided some explanation of the increasing income-child health gradient. One possible mechanism of this increasing trend found by Currie and Stabile was that low-SES Canadian children were subject to frequent health shocks. Using a regional UK birth cohort data Propper, Rigg, and Burgess (2007) revealed that mother's health, particularly her mental health, could explain the relationship between child health and family income in the UK. They found no direct effect of income if parental health and parental child health related behaviour are controlled for. Khanam et al. (2009) also found that parental health, in particular, mother's health could explain some of this mechanism in their Australian sample. These two studies from the UK and Australia, that used sample of children with similar age (0-7 years), found that the relationship between income and child health disappears if parental health is considered. Khanam et al. (2009), however, did not look at the two other possible mechanisms of increasing income gradient proposed by Currie and Stabile (2003).

First hypothesis of Currie and Stabile (2003: p1813) is that "low-SES children are less able to respond to a given health shock, so that the negative effects of health shocks persist and accumulate over time". The second hypothesis is "low-SES children respond to health shocks in a way that is similar to high-SES children, but are subject to more health shocks". J. Currie and Stabile (2003) found that the increasing gradient in child health in Canada was not because of these children lack resources to respond to a health shock. They found that low-SES children responded to a particular health shock to a similar rate as high SES children as they age, but they were subject to new health shocks. However, a recent study from the USA by Condliffe and Link (2008, subsequently, C&L) showed that low-SES children responded differently to past health shock compared to high-SES children. They also found some evidence that low-SES children were subject to more health shocks as they age, however, this effect was not as strong as J. Currie and Stabile's study. Wei (2007)'s study, using the 1996-2004 Medical Expenditure Panel Survey (MEPS), also confirmed that children from high-income

¹Case et al used National Health Interview survey (NHIS) of America

USA households recovered better from a past health shocks compared to children from low-income household. However, she found that children from low-income family were not subject to higher incidence of health shock, rather, their parents were less able to manage a health shocks, which in turn, left these children in poor health. Murasko (2008) focussed on the different dimension of the increasing gradient. He found that baseline health (i.e., previous health state), which was described as 'cumulative effect of income' in his paper, could explain the increasing income-child gradient, at least until they reached adolescent. These confronting results from the USA and Canada make a compelling case to explore this issue further by exploiting longitudinal data from a different country.

In this paper, we explain the steepening gradient in child health in Australia using panel data from the LSAC, focusing mainly on the two hypotheses proposed by Currie and Stabile. We will also test the 'cumulative effect' and 'contemporaneous effect' of income hypothesis proposed by Murasko (2008).

2 Literature Review

As we discussed in the previous section, Currie and Stabile (2003) found that the increasing income gradient in child health in Canada was not because low-SES children were affected most by a health shock, rather they experienced more health shocks compared to high-SES children. A recent study from the USA by Condliffe and Link (2008), however, revealed that low-SES children were hit harder by a health shock because of lower parental resources to respond to a health shock. They also found that low-SES children were more likely to experience the incidence of chronic condition as they age, which could explain why income-health gradient was stronger for older children. Interestingly, the income gradient found in Condliffe and Link (2008), using the 1996-2002 Medical Expenditure Panel Survey (MEPS), was flatter than Case et al (2002). Surprisingly, the gradient found in their study for the USA was even lower than Canada (Currie and Stabile, 2003), which has a universal health insurance scheme.

Wei (2007) examined the role of chronic condition in understanding the increasing income-child health gradient using the 1996-2004 MEPS from the USA. She found that child health was affected by both current and past shocks. She also found that children from low-income family were not subject to more health shock rather their parents were less able to manage a health shock, which in turn, left these children in poor health. Her findings also confirmed that children from high-income USA households recovered better from a past health shock than the low-income household. She defined this as 'disease management ability' of parents. Moreover, the author revealed that family income was not protective in reducing morbidity in children who were having mild or time intensive chronic condition. While children with severe and financially intensive condition from high-income households recovered better from the shock compared to low-income households with the same condition. She argued that if parents were more capable to manage a chronic disease, they could protect their children from having a bad health. However, parental disease management ability could be a function of income, which can explain the stronger relationship between income and health in children with chronic condition.

Using the 1996-2005 MEPS from the USA, Murasko (2008) found an increasing income-child health gradient as Case et al. (2002), however, this association (smaller coefficients on income over the different age group) in MEPS was much weaker than the 1995 NHIS used by Case et al. He argued that it was not possible in the cross section (e.g., Case, Lubotsky, and Paxson, 2002) whether the increasing income-child health association across age-groups was due to a cumulative effect, a contemporaneous effect, or a combination of both. He proposed to consider baseline health (health

states of a child at the beginning of the survey) in the model to account for cumulative effect (past influence) of income on health. He explained the coefficient of income as a contemporaneous effect and coefficient of baseline health as a cumulative effect of income. He found that if the baseline health was considered in the model, the child health-income gradient became flattened over the age group, however poor health was still more prevalent in older children. Therefore, he concluded that it was the cumulative effect that could explain much of the increasing gradient in child health and income, particularly, before adolescence, while a strong contemporaneous association remained for this age group.

In this paper, we propose that 'cumulative effect' of income can also be explained by a past health shock (in the form of a chronic condition or an injury resulting from an accident), while income coefficient explains the 'contemporaneous effect'.

The above-mentioned papers emphasise on the role of chronic condition in the explanation of income and child health gradient. The studies from the USA and Canada are vivid in this debate. Murasko (2008) argued that poor health is more prevalent in older children compared to younger children, and this can explain some of the steepening gradient in child health in the USA.

3 Data and Descriptive Statistics

This study utilises the data from the first two waves of the nationally representative Longitudinal Study of Australian Children (LSAC) (Australian Institute of Family Studies, 2007). The LSAC has so far involved two waves of data collection for more than ten thousand children. The LSAC collects data on these children every two years and will follow them until 2010 or beyond. The LSAC was conducted using both face-to-face interviews and survey instruments that were sent and retrieved via mail. The main topics covered include demographics, health status, education, the relationship history of parents, parenting practices, financial factors, lifestyle, housing and neighbourhood attributes.² The data were collected using a two-stage clustered sampling design with postcodes were used as the primary sampling unit (PSU). To ensure proportional geographic representation, postcodes were selected as a stratified sample by state of residence, and urban and rural geographical status. The sampling frame for the second stage consisted of all children born in the selected PSUs between March 2003 and February 2004 (B-Cohort, infants aged 0-1 years in 2004), and between March 1999 and February 2000 (K-Cohort, children aged 4-5 years in 2004) who were enrolled on the Health Insurance Commission's Medicare database. The Australian Medicare scheme is universal and compulsory; thus the sample constructed for the LSAC is generally representative of Australian children in these age cohorts, although children living in remote areas were not sampled.

The LSAC approach results in a sample frame that contains approximately 5000 children in each cohort, with an average of 20 children per cohort per postcode. The final respondent samples consist of 5107 and 4983 children in cohorts B and K, respectively, in Wave 1 (conducted in 2004). The numbers of children surveyed in Wave 2 (conducted in 2006) of the respective cohorts is slightly lower, primarily as a result of attrition, with 4606 and 4464 children retained in cohorts B (aged 2-3 years in 2006) and K (aged 6-7 years in 2006), respectively. The attrition rates are therefore 9.8 and 10.4 per cent for B and K cohorts, respectively. The logistic regressions conducted by Mission and Siphthorp (2007, Tables 1-2) reveal that attrition occurred mostly at random in the LSAC. However, attrition was slightly more likely if Parent 1 (primary caregiver) is a young male, the household was living in a

²For a more comprehensive account of the LSAC sampling frame of the LSAC see Soloff, Lawrence, and Johnstone (2005).

Table 1: Descriptive statistics

| Description | Obs | Means | Linearised Std. Err. | Min | Max |
|-------------------------------------|-------|-------|-------------------------|-------|-------|
| Log of income | 18929 | 10.99 | 0.02 | 3.26 | 13.24 |
| Child's age (months) | 18929 | 82.09 | 0.05 | 3.00 | 94.00 |
| Gender of the child (male=1) | 18929 | 0.51 | 0.01 | 0.00 | 1.00 |
| Log of household size | 18929 | 1.49 | 0.00 | 0.69 | 2.64 |
| Both biological parents present | 18929 | 0.80 | 0.01 | 0.00 | 1.00 |
| Mother age at birth (years) | 18833 | 29.67 | 0.11 | 13.00 | 63.00 |
| Mother completed year 12 | 18929 | 0.49 | 0.01 | 0.00 | 1.00 |
| Mother had a graduate degree | 18929 | 0.19 | 0.01 | 0.00 | 1.00 |
| Mother had a postgraduate degree | 18929 | 0.05 | 0.00 | 0.00 | 1.00 |
| Primary caregiver is not mother | 18929 | 0.10 | 0.01 | 0.00 | 1.00 |
| Low income household* | 18929 | 0.11 | 0.01 | 0.00 | 1.00 |
| General health status in Wave 1 | 9009 | 1.59 | 0.01 | 1.00 | 5.00 |
| Poor health of the child** | 18929 | 0.13 | 0.01 | 0.00 | 1.00 |
| Number of chronic conditions | 18929 | 0.36 | 0.01 | 0.00 | 7.00 |
| Number of conditions in Wave 1 | 18929 | 0.28 | 0.00 | 0.00 | 7.00 |
| Number of new chronic conditions*** | 18929 | 0.18 | 0.01 | 0.00 | 5.00 |
| Having new health shock | 18929 | 0.16 | 0.01 | 0.00 | 1.00 |
| Chronic condition in Wave 1 | 9010 | 0.34 | 0.01 | 0.00 | 1.00 |
| Asthma in Wave 1 | 4420 | 0.21 | 0.01 | 0.00 | 1.00 |
| Injury in Wave 1 | 9010 | 0.17 | 0.01 | 0.00 | 1.00 |

Note: * Determined using the poverty line of Australia (Melbourne Institute, 2008); **health status good, fair and poor, ***Number of chronic conditions appear in Wave 2 but not in Wave1; Means and standard errors are estimated using the Taylor linearisation method (Kish, 1995; Chambers and Skinner, 2003)

Source: Computed from the Longitudinal Study of Australian Children (Australian Institute of Family Studies, 2007).

rented home, or in an areas with a lower socio-economic status index. For the B-cohort, attrition was also more likely to occur among households in areas where fewer people in the postcode speak only English in the home.

In order to take the advantage of the survey's design characteristics, all analyses presented in this paper apply the sampling weights of the LSAC. These are computed as the inverse of the probability of a child being selected for inclusion in the LSAC sample. For example, if the probability of a child is being sampled is 0.20, the weight given to that child's response is 5.0. In addition, cluster information are used to produce correct variances of the estimates as there is less variations among variables within a cluster (i.e., postcode). This approach also corrects for the fact that the variance is reduced in a finite population with non-replacement sampling (i.e., in non-replacement samples, the population being sampled is reduced as the sampling progresses; and the variance is thereby reduced). The descriptive statistics of the variables are reported in Table 1.

3.1 Child Health

As with the foregoing literature on income and child health (see for example, Case et al., 2002; J. Currie and Stabile, 2003; A. Currie et al., 2007), our measure of child health is constructed from the following question that was asked to the child's primary care-giver (Parent 1)³: "*In general, how would you say child's current health is?*" The responses were recorded on a five-point Likert scale upon which 1 is "Excellent" 2 is "Very good"; 3 is "Good"; 4 is "Fair" and 5 is "Poor". Other researchers

³In principle, Parent 1 is the person in the family who knows the most about the study child. In most cases this is the child's biological mother but, alternatively may be the biological father, a step-parent, an adoptive parent, a guardian, or someone else who has a parental/guardian relationship with the child.

have found that there are typically very few respondents in the “Poor” health category: in the LSAC approximately 0.30 per cent of the children sampled fall into this category. Some authors (e.g., A. Currie et al., 2007) have chosen to merge the lowest and second-lowest health state categories as a response to the (relatively) small number of observations in the “Poor” health category. Since there are no shortage of degrees of freedom in our study, we do not compress the “Fair” and “Poor” categories of child health. Thus, our dependent variable for parent-reported overall child health contains the five original categories.⁴

3.2 Chronic Condition

The diagnosis of chronic condition is a function of child’s age, as the children in our sample is young compared to the previous studies, therefore the prevalence of chronic condition is likely to be lower in our sample. In the LSAC, Parent 1 was asked whether or not the child had a long-term medical condition, the nature of the condition and whether the child had experienced any developmental delays that were attributable to the problem, compared to children of a similar age. If the answer was yes, the respondent was asked to check up to fourteen chronic conditions. Approximately thirty per cent of survey children in the LSAC were reported to have at least one such condition, and seven per cent have more than one such condition.⁵ Furthermore, the LSAC contains information on whether the child has asthma or bronchiolitis, *as diagnosed by a health professional*.⁶ The survey revealed that 19.32 and 13.32 per cent of children, respectively, were reported to have been diagnosed with asthma and bronchiolitis (see Table 2). Approximately 9 per cent of children, who did not have a condition in wave 1, have been diagnosed with a new chronic condition in wave 2. The LSAC revealed that 12.58%, 30.48% and 20.82% children respectively were reported as having poor health (poor health = 1 if a child is reported as having good, fair and poor health category), suffering from a chronic condition and asthma in wave 1.

We are interested in examining how the parents manage a previous health shock that affects current health status of a child. We also investigate factors determining whether or not a child had an injury in the wave 1.⁷ We assume that injury is a health shock that could affect health. The LSAC reported that 11.63% children had at least one injury in Wave 1 whilst the figure of Wave 2 is 17.66%.

4 Method

Following Grossman (1972) we argue that a child is born with a initial stock of health that depreciates over time. The arrival and severity of different health shocks, and insufficient investment into health capital could accelerate the rate of depreciation of the health stock. However, health stock should be retained to a level that is above to a critical minimum level for survival. This conceptual framework could explain the increasing gradient by child age. Suppose, if the depreciation rate of health stock of older children is higher than the younger children, then that will result in an increasing gradient over the age-group. As a child grows older, he is susceptible to more health shocks, in the form of chronic

⁴Nevertheless, we also conducted analysis with the last two categories recoded and the results show little differences. These estimates are available from the authors upon request.

⁵Chronic conditions include hearing problems, vision problems,eczema, diarrhea/collitis, ear infections, other infections, food or digestive allergies, other illness, other physical disabilities, recurrent abdominal pain, development delay, anaemia, attention deficit disorder and frequent headaches.

⁶The survey questions for this variable is “Has a doctor ever told you that you child has: asthma?, bronchiolitis?”

⁷The primary care-giver was asked: “what types of injury or accident did child have that needed medical attention? Ten types of injuries were reported: broken or fractured bones, burn or scald, dislocation, sprain or strain, cut or scrape, concussion or internal head injury, internal injury (not head), dental injury, accidental poisoning and other.

Table 2: Children with chronic conditions and injury by SES and Wave

| Waves | SES | Any chronic condition | Asthma | Injury |
|--------|------|-----------------------|--------|--------|
| Wave 1 | Low | 31.24% | 22.09% | 12.29% |
| | High | 30.20% | 20.39% | 11.34% |
| | All | 30.48% | 20.82% | 11.63% |
| Wave 2 | Low | 28.40% | 20.49% | 17.75% |
| | High | 29.89% | 17.05% | 16.51% |
| | All | 29.41% | 18.51% | 17.66% |
| Both | Low | 30.02% | 21.13% | 14.64% |
| | High | 30.05% | 18.26% | 13.87% |
| | All | 29.97% | 19.32% | 14.50% |

Note: The definition of low SES and high SES was applied according to Condliffe and Link (2008). Low SES includes those whose income is below the mean income by one standard deviation while high SES include those whose income is above the mean level by one standard deviation.

Source: Computed from the Longitudinal Study of Australian Children (Australian Institute of Family Studies, 2007).

condition, injury from an accident, which could depreciate the health stock. If proper investment is not made to recover from those health shock then health disparities will increase. The depreciation rate might be higher for children from low-SES, and children with chronic condition. Because, children from low-SES households are more likely to have a health shock (because of differences in lifestyle, poor housing quality, inadequate nutrition) and recover slowly from a health shock because of lack of information or resource constraint of the parents. Low-SES parents are less able to manage a health shock as effectively as high-SES parents, which leaves their children in poor health. These propositions could explain the increasing income-child health gradient. Therefore, family income could affect child health through the incidence of chronic condition and different recovery rate because of differential investment to the health stock.

Previous research has identified two possible channels by which income gradient may increase by child age. First, children of low-SES parents may respond to health shock differently than the high-SES parents. That is low-socio-economic status children may recover slowly from a particular health shock than the high-SES children, so negative effects of health shock accumulate over time, and may contribute to the increasing gradient. Second, Low-SES children may experience more incidence of health shock (chronic conditions, diseases requiring medical attention, different forms of injuries). Therefore, the depreciation rate of health stock of low SES children could be higher than the high SES children as a results of these two channels. The recent literature (Condliffe and Link, 2008) has established that differential response to past health shocks by SES can contribute to the increasing gradient. Children of low-SES parent are also more likely to have more health shock than the high-socioeconomic status.

The following empirical approach is adopted to explore these issues in a multivariate context. We first proceed by exploring the income gradient in child health. In an attempt to do this, we pool data from the first two waves from the LSAC to replicate the results of Currie and Stabile (2003) and Condliffe and Link (2008). We estimate the following equation:

$$health_{it} = \alpha_0 + \alpha_1 \ln(inc)_{it} + \alpha_2 age_{it} + \alpha_3 X_{it} + \varepsilon_{it} \quad (1)$$

where, $health_{it}$ is a child's health status (in time t) measured on 1-5 point Likert scale with 1 indicating excellent and 5 indicating poor health. The term, $\ln(inc)_{it}$ is log of permanent family income of a household which is believed to have stronger effect on health than transitory variations in income.

We take the average annual income of each family in Wave 1 and Wave 2. The income estimates were then adjusted using the Australian national Consumer Price Index (CPI) for the study period, using the CPI at Wave 1 as the base (Australian Bureau of Statistics, 2008). The term age_{it} is a set of age dummies, and X_{it} is a set of exogenous variables that include a set of dummies for mother education, wave dummies, cohort dummies, log of family size, a dummy variable for the sex of the child, a dummy variable for having a primary care giver (*person responding to the survey questions*) that is not the biological mother, a dummy variable for having a female care giver, a dummy variable for having two biological parents in the household, and the mother's age at birth. As the children in our sample are for ages 0-8 years, so we estimate equation (1) for the age groups: 0-3years and 4-8 years. The equation 1 is a basic equation used in the previous literature (for example, Case, Lubotsky, and Paxson, 2002; Currie and Stabile, 2003; Condliffe and Link, 2008; Murasko, 2008) to examine the relationship between child health and household income. Given the ordered nature of $health_{it}$, equation 1 is estimated by a ordered probit model. We expect a negative sign for α_1 and the absolute value of α_1 will increase over the age groups suggesting an increasing income-child health gradient.

Murasko (2008) argued that it is not possible in cross section (for example, eq.1) whether any increasing income-child gradient is due to a cumulative, a contemporaneous effect, or a combination of both. The cumulative effect of income is consistent with the idea that prevalence of poor health in low-SES children in their early years of life might be carried out into their adulthood, which in turn increases health disparities among low and high SES. This could be one reason why income gradient increases by age. The contemporaneous effect suggests that the health disadvantages faced by the low-SES households do not become obvious until the children grow older. He proposed to add baseline health in eq (1) to evaluate these potential effects.

Therefore, to test the 'cumulative' and 'contemporaneous' effect of income, we propose the following model for child health:

$$health_{it} = \beta_0 + \beta_1 \ln(inc)_{it} + \beta_2 age_{it} + \beta_3 health_{it-1} + \beta_4 X_{it} + \varepsilon_{it} \quad (2)$$

where $health_{it-1}$ is child's general health state measured in time $t - 1$, which is wave 1 in the LSAC; the remaining variables are defined as above.

Now we proceed to test the two hypothesis proposed by Currie and Stabile (2003) and evidenced by Condliffe and Link (2008) by exploiting panel data of LSAC. We estimate the following equation that is basically based on Currie and Stabile (2003) and Condliffe and Link (2008):

$$health_{it} = \gamma_0 + \gamma_1 \ln(inc)_{it} + \gamma_2 age_{it} + \gamma_3 shock_{i(t-1)} + \gamma_4 \ln(inc)_{it} * shock_{it-1} + \gamma_5 X_{it} + \varepsilon_{it} \quad (3)$$

where, $health_{it}$ is a measure of child health in wave 2; $shock_{i(t-1)}$ is a health shock in wave 1; and rest of the variables are defined as above. The variable $health_{it}$ is a dichotomous variable equals to one if health status is poor, fair, or good; $shock_{i(t-1)}$ is also a dichotomous variable which equals 1 if the child has any chronic condition, asthma or an injury in wave 1. The equation 3 is measured by binary probit model. A statically significant and negative coefficient for γ_4 would indicate that low-SES children will be severely affected by a health shock.

The second hypothesis is tested by the following equation:

Table 3: The increasing income gradient in child health (ordered probit models)

| Variables | 0-3 years old | | | 4-8 years olds | | |
|-------------------------------|-----------------------|----------------------|-------------------------|-----------------------|----------------------|-------------------------|
| | Canada (C&S, 2003) | USA (C&L, 2008) | Australia This paper | Canada (C&S, 2003) | USA (C&L, 2008) | Australia This paper |
| Without mother's education | ***-0.151 (0.026) | ***-0.136 (0.018) | ***-0.078 (0.027) | ***-0.216 (0.019) | ***-0.204 (0.014) | ***-0.148 (0.023) |
| With mother's education | ***-0.132 (0.027) | ***-0.093 (0.018) | ***-0.083 (0.027) | ***-0.182 (0.020) | ***-0.156 (0.014) | ***-0.144 (0.025) |

Notes: (i) The dependent variable is an ordered categorisation of the child's general health status (e.g., 1 = excellent, 2 = very good, 3 = good, 4 = fair and 5 = poor) as reported by a parent/guardian. (ii) As the LSAC data are only available for children aged 0-8, we report the results for same age groups from previous studies, though those studies also included children older than 8 years. (iii) Robust standard errors are in parentheses. (iv) ***Significant at the one per cent level.

Sources: J. Currie and Stabile (2003), Condliffe and Link (2008). Australian estimates were computed from the Longitudinal Study of Australian Children (Australian Institute of Family Studies, 2007).

$$Newshock_{it} = \delta_0 + \delta_1 In(inc)_{it} + \delta_2 age_{it} + \delta_3 In(inc)_{it} * age_{it} + \delta_4 shock_{t-1} + \delta_5 X_{it} + \varepsilon_{it} \quad (4)$$

where, $Newshock_{it}$ refers to number of new conditions in wave 2 and other variables are defined as above. The dependent variable, $Newshock_{it}$, equals 1 if a child develops a new condition between two periods. The equation 4 is estimated by a binary probit model. We will estimate the several forms of the above equation to test the hypothesis: children from low-SES experience more incidence of health shock.

5 Results

5.1 The Gradient:Cross-Sectional Estimates

The equation 1 is estimated for the children aged 0-3 years and aged 4-8 years, with and without mother education to compare with C&S and C&L. Our results from the LSAC and C&S and C&L results are reported in Table 2. It is noteworthy that C&S's and C&L's analysis were for the children aged 0-15 year and aged 0-17 year respectively. As LSAC provides data for the children 0-8 years, so we will replicate the results for these age groups from C&S and C&L. In line with the previous literature, the results from the LSAC show an increasing income and child health gradient in Australia for 0-8 years. However, the association between family income and child health is weaker in the LSAC data (the coefficient on income in Australia is lower than the Canada and the USA). The inclusion of mother education has no effect on the magnitude of the coefficients on income in Australia, while, inclusion of maternal education reduced the magnitude of income coefficient in Canada and in the USA.

5.2 Causes of the Gradient:Panel Estimates

In this section we explore the underlying mechanisms of the increasing gradient exploiting the panel data of LSAC. A recent study by Murasko (2008; p 1490) argued that it is not possible in a cross-sectional analysis whether a steepening gradient is because of 'a cumulative effect of income on health through childhood and adolescence or whether it reflects a stronger contemporaneous effect of income on health for older children. To find out whether there is any cumulative effect and or contempora-

Table 4: The effects of based line health on current health status.

| Variables | 0-3 years | | 4-8 years | |
|--|---------------------------------|-------------------------|--------------------------------|-------------------------|
| | United States Murasko (2008) | Australia This Paper | United States Murasko(2008) | Australia This Paper |
| Ordered probit estimates | | | | |
| Log of income | -0.025 (0.021) | -0.066 (0.051) | ***-0.059 (0.021) | ***-0.123 (0.034) |
| Health _{<i>i</i>-1} =1 (excellent) | ***0.523 (0.040) | -0.276 (0.406) | ***0.585 (0.036) | ***-2.140 (0.501) |
| Health _{<i>i</i>-1} =2 (very good) | ***0.851 (0.043) | 0.182 (0.404) | ***0.962 (0.048) | ***-1.490 (0.502) |
| Health _{<i>i</i>-1} =3 (good) | ***1.335 (0.095) | 0.500 (0.410) | ***1.368 (0.087) | ** -1.076 (0.504) |
| Health _{<i>i</i>-1} =4 (fair) | | 0.644 (0.424) | | -0.758 (0.530) |
| N | 7596 | 3323 | 10352 | 4520 |
| Probit estimates | | | | |
| Log of income (lninc) | -0.010 (0.007) | 0.023 (0.068) | ** -0.012 (0.006) | ***-0.151 (0.048) |
| Health _{<i>i</i>-1} =3,4 or 5 (good, fair or poor) | ***0.229 (0.016) | ***0.727 (0.068) | ***0.279 (0.017) | ***0.989 (0.060) |
| N | 7596 | 3315 | 10352 | 4520 |

Note: (i) Robust standard errors are in parentheses. (ii) Significant levels are ***=1%, **=5%, and *=10%

Sources: Condliffe and Link (2008)). Australian estimates were computed from the Longitudinal Study of Australian Children (Australian Institute of Family Studies, 2007).

neous effect of income in the steepening income-child health gradient, and whether or not the chronic condition plays any role in explaining the gradient we exploit panel data from the LSAC. First, we focus on Murasko(2008)'s 'cumulative effect and contemporaneous effect of income', then we test the hypotheses of C&S.

5.2.1 Cumulative Effect Versus Contemporaneous Effect

We estimate eq (2) to examine whether the cumulative effect and contemporaneous effect of income can explain the increasing gradient in the LSAC data. Our other covariates are similar to C&S to keep consistency with rest of the paper. However, we also estimate eq(1) and eq (2) using similar variables as Murasko. The results of these specifications(not reported in the paper) show similar trends. When we control for baseline health status of a child, the income coefficient becomes insignificant for the 0-3 age group. Murasko's also found insignificant coefficient on income for this age group in the USA. Murasko (2008) revealed that controlling for the baseline health status flattens the income gradient in the USA, our results also show similar trends over the age groups: 0-3 years and 4-8 years.

Murasko argued that if cumulative effect is dominant, then the coefficient on income (α_1) would be relatively flatter. If contemporaneous effect is dominant in the steepening income gradient, then the coefficient on income (α_1) would be stronger over the age group.. In line with Murasko, if we explain previous health states as a 'cumulative effect' of income, we find that controlling for baseline health (cumulative effect) makes the income-child health gradient flatter. This findings from the LSAC suggest that 'cumulative effect' can explain some of the increasing gradient in income and child health. The incidence of poor health increases over the age groups, which is also consistent with Murasko's finding. The high prevalence of poor health among the 4-8 years may partly explain why the relationship between income and child health becomes stronger for the older children.

5.2.2 The Role of Chronic Condition

In this section, we follow J. Currie and Stabile (2003) and Condliffe and Link (2008) to test the hypotheses originally proposed by J. Currie and Stabile.

Is There any Differential Response by Different SES to Past Health Shocks?

In this section we examine whether low-SES children are hit harder by a health shock. Following C&S and C&L we estimate different versions of Equation 3 by binary probit models. We report the results of J. Currie and Stabile (2003), Condliffe and Link (2008) and this paper in Table 5.

Our results show that earlier health shock increases the probability of having poor health in current period. For example, the presence of chronic conditions in Wave 1 (2004) increases the probability of being in poor health in wave 2 (2006) by 7 percentage points (estimated from relevant marginal effects); the estimated coefficient is statistically significant at the 1 percent level. The positive and statistically significant (at the 1 percent level) coefficient on asthma show that having asthma in Wave 1 (in 2004) increases the likelihood of being reported in poor health in Wave 2 (in 2006) by 10 percentage points (estimated from corresponding marginal effects). However, the coefficients of earlier shocks become statistically insignificant when an interaction term between health shock and family income is used in the regression. It is noteworthy that having an injury in Wave 1 does not increase the probability of poor health in Wave 2.

The coefficients on income are always negative and statistically significant at the 1 % level in these specifications indicating a protective effect of income on child health. However, the interaction term between family income and a lagged health condition is not significant and in some cases the sign is positive. This result is not surprising because children in our sample are younger (0-7 years) than those of J. Currie and Stabile and Condliffe and Link. These children are perhaps too young to capture any income related disparity. However, the negative and statistically insignificant sign for the interaction terms indicate that the low-SES household do not respond to a health shock differently compared to high-SES household. This finding is in line with J. Currie and Stabile (2003) but in contrast to Condliffe and Link (2008). Therefore, our results do not provide any evidence that low-SES households respond to a past health shock differently than the high-SES households or they are less able to manage a health shock than the high-SES household.

New Health Shock

We estimate different forms of Equation (4) by probit models to see whether a child from low-SES background is more likely to experience a new health shock. First we include income with the standard control variables, then we include an interaction term of income and age. Finally, we include an indicator for past health shock that equals 1 if the child had any chronic condition or asthma in wave 1. Our results indicate that SES plays no role on the probability of having a new shock. Our results are in line with Wei (2007)'s who also found that children from low income household are not subject to new health shock. Although C&S established that low-SES Canadian children were subject to more health shocks, subsequent studies from the USA do not find any conclusive evidence in support of this. For example, C&L used two datasets from the USA, in one data set (PSID) they found no evidence that low-SES children are more likely to experience new health shock.

Our results reveal that children who experienced past healths are more likely to experience new health shocks. For example, children who had a health shock in wave 1 (in 2004) are 4 percentage points more likely to have a new health shock in wave 2 (in 2006). The children who had asthma in

Table 5: The effects of earlier health conditions on poor health today

| <i>Canadian evidence (J. Currie and Stabile, 2003)</i> | | | | | | |
|---|-----------------------|-----------------------|-----------------------|-----------------------|----------------------|-----------------------|
| Variables | (1) | (2) | (3) | (4) | | |
| Chronic Condition in 1994 | **0.108 (0.008) | *0.257 (0.138) | | | | |
| Asthma in 1994 | | | **0.136 (0.012) | *0.357 (0.209) | | |
| Log of average income | **_-0.055 (0.007) | **_-0.052 (0.007) | **_-0.055 (0.007) | **_-0.053 (0.007) | | |
| <i>Interactions of the logs average income with</i> | | | | | | |
| Chronic Condition in 1994 | | -0.014 (0.013) | | | | |
| Asthma in 1994 | | | | -0.021 (0.019) | | |
| <i>American evidence (Condliffe and Link, 2008)</i> | | | | | | |
| Variables | (1) | (2) | (3) | (4) | | |
| Chronic Condition in 1997 | **0.0567 (0.024) | **0.631 (0.303) | | | | |
| Asthma in 1997 | | | **0.086 (0.036) | 0.549 (0.418) | | |
| ln (family income) | **_-0.055 (0.019) | **_-0.040 (0.020) | **_-0.054 (0.019) | **_-0.048 (0.012) | | |
| <i>Interactions of the logs average income with:</i> | | | | | | |
| Chronic Condition in 1997 | | **_-0.054 (0.028) | | | | |
| Asthma in 1997 | | | | -0.044 (0.039) | | |
| <i>Australian evidence (This Paper)</i> | | | | | | |
| Variables | (1) | (2) | (3) | (4) | (5) | (6) |
| Log of income | ***_-0.143 (0.030) | ***_-0.140 (0.036) | ***_-0.245 (0.043) | ***_-0.225 (0.047) | ***_-0.141 (0.03) | ***_-0.148 (0.031) |
| Chronic Condition in Wave 1 (2004) | ***0.307 (0.035) | 0.414 (0.623) | | | | |
| Asthma in Wave 1 (2004) | | | ***0.431 (0.060) | 1.319 (0.938) | | |
| Injury in Wave 1 (2004) | | | | | 0.028 (0.053) | -0.651 (0.845) |
| <i>Interactions of the logs of average income with:</i> | | | | | | |
| Chronic Condition in Wave 1 | | -0.010 (0.056) | | | | |
| Asthma in Wave 1 | | | | -0.081 (0.085) | | |
| Injury in Wave 1 | | | | | | 0.062 (0.076) |
| N | 8957 | 8957 | 4381 | 4381 | 8957 | 8957 |

Notes: (i) Robust standard errors are in parentheses. (ii) Significant levels are ***= 1%, **=5%, and *=10% (iii) For details of the Specifications for the J. Currie and Stabile (2003) see their Table 3 and for the Condliffe and Link, (2008) see their Table 8. (iv) Australian estimates are computed from the Longitudinal Study of Australian Children (Australian Institute of Family Studies, 2007).

Table 6: The effects of income and past health shocks on new health shocks

| Variables | (1) | (2) | (3) | (4) |
|--------------------------------------|------------------|--------------------|---------------------|---------------------|
| Log of family income | 0.022 (0.025) | 0.032 (0.054) | 0.029 (0.054) | 0.299 (1.006) |
| Interaction of age and income | | -0.0001 (0.001) | -0.0001 (0.001) | -0.004 (0.012) |
| Chronic condition in previous period | | | ***0.124 (0.031) | |
| Asthma in the previous period | | | | ***0.180 (0.050) |
| N | 8957 | 8957 | 8957 | 4381 |

Notes: (i) Robust standard errors are in parentheses. (ii) Significant levels are ***=1%, **=5%, and *=10%

Table 7: Income gradient with control for health expenditure

| Variables | Health status | | Health expenditure | |
|---------------------------------|---------------|-----------|--------------------|-----------|
| | coef. | std. err. | coef. | std. err. |
| Log of income | ***-0.108 | 0.011 | -0.020 | 0.020 |
| Gender of the child | ***0.112 | 0.020 | ***0.148 | 0.023 |
| Log of hh size | **0.085 | 0.036 | ***-0.44 | 0.047 |
| Both biological parent presence | -0.039 | 0.033 | *0.076 | 0.040 |
| Mother's age at birth | ***-0.006 | 0.002 | **0.005 | 0.002 |
| Mother's education > year 12 | -0.003 | 0.038 | 0.009 | 0.025 |
| First caregiver is not mother | ***-0.129 | 0.016 | ***0.685 | 0.024 |

Note: Wald test for independent of equations, $Chi^2(1)=963.15$, p -value=0.00. Significant levels are ***=1%, **=5%, and *=10%

wave 1 (in 2004) are 5 percentage points more likely to diagnose with a new health condition in wave 2 (in 2006).⁸

6 Conclusions

Our results show that none of the hypotheses proposed by C&S can explain the increasing income-child health gradient in Australia. Our results revealed that low-SES Australian children are not severely affected by a health shock, and they are not subject to more health shock as they age. Our results find support for Murasko(2008)'s cumulative effects of income hypothesis. Our results show that the incidence of poor health increases by child age, which indicates that cumulative effects of baseline health contribute to poor health of the older children. However, it has some role to reduce the increasing gradient, but it does not make the income coefficient insignificant. It is noteworthy that our analysis is for the children aged 0-7 years, while the literature in this field is for the children aged 0-15/17 years. Therefore, our analysis might not be easily comparable to the whole range of findings from the existing literature. However, it is still comparable to the same age group and provides some important insights that could be interesting for policy purpose. So far, from the existing literature, we do not see any uniform mechanism that can be applied universally. So, policies to address this issue should be country specific.

⁸These figures - not reported in Table 5 - are obtained from the corresponding marginal effects .

References

- AUSTRALIAN BUREAU OF STATISTICS (2008): “Consumer Price Index, Australia, Jun 2008,” Australian Bureau of Statistics <http://www.abs.gov.au/Ausstats/abs@.nsf/mf/6401.0>, Assessed 1.7.2008.
- AUSTRALIAN INSTITUTE OF FAMILY STUDIES (2007): “Longitudinal Study of Australian Children, Wave 2 Data Release,” Australian Institute of Family Studies, Melbourne, www.aifs.gov.au/growingup.
- CASE, A., D. LUBOTSKY, AND C. PAXSON (2002): “Economic status and health in childhood: The origins of the gradient,” *The American Economic Review*, 92(5), 1308–1344.
- CONDLIFFE, S., AND C. R. LINK (2008): “The Relationship between Economic Status and Child Health: Evidence from the United States,” *American Economic Review*, 98(4), 1605–1618.
- CURRIE, A., M. A. SHIELDS, AND S. W. PRICE (2007): “The child health/family income gradient: Evidence from England,” *Journal of Health Economics*, 26(2), 213–232.
- CURRIE, J., AND M. STABILE (2003): “Socioeconomic Status and Child Health: Why Is the Relationship Stronger for Older Children,” *The American Economic Review*, 93(5), 1813–1823.
- GROSSMAN, M. (1972): “On the Concept of Health Capital and the Demand for Health,” *Journal of Political Economy*, 82(2), 223–255.
- KHANAM, R., H. S. NGHIEM, AND L. B. CONNELLY (2009): “Child Health and the Income Gradient: Evidence from Australia,” *Journal of Health Economics*, (Forthcoming).
- MISSION, S., AND M. SIPTHORP (2007): “LSAC Technical Paper No.5: Wave 2 weighting and non-response,” Discussion paper, Australian Institute of Family Studies.
- MURASKO, J. E. (2008): “An evaluation of the age-profile in the relationship between household income and the health of children in the United States.,” *Journal of Health Economics*, 27(6), 1407–1652.
- PROPPER, C., J. RIGG, AND S. BURGESS (2007): “Child health: evidence on the roles of family income and maternal mental health from a UK birth cohort,” *Health Economics*, 16(11), 1245–1269.
- SOLOFF, C., D. LAWRENCE, AND R. JOHNSTONE (2005): “LSAC Technical Paper No. 1: Sample design,” Discussion paper, Australian Institute of Family Studies, Melbourne, Australia.
- WEI, D. (2007): “Family Income and Child Health: The Role of Chronic Conditions,” Discussion paper, Department of Economics, University of Wisconsin Madison.