

**Rapid-Cycle PCR Method To Detect
Bordetella pertussis That Fulfills All
Consensus Recommendations for Use of
PCR in Diagnosis of Pertussis**

D. J. Farrell, M. McKeon, G. Daggard, M. J. Loeffelholz, C. J. Thompson and T. K. S. Mukkur
J. Clin. Microbiol. 2000, 38(12):4499.

Updated information and services can be found at:
<http://jcm.asm.org/content/38/12/4499>

These include:

REFERENCES

This article cites 14 articles, 13 of which can be accessed free at: <http://jcm.asm.org/content/38/12/4499#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Rapid-Cycle PCR Method To Detect *Bordetella pertussis* That Fulfills All Consensus Recommendations for Use of PCR in Diagnosis of Pertussis

D. J. FARRELL,^{1,2*} M. McKEON,² G. DAGGARD,² M. J. LOEFFELHOLZ,³
C. J. THOMPSON,³ AND T. K. S. MUKKUR²

Microbiology Department, Toowoomba Laboratory, Queensland Health Pathology Service,¹ and Centres for Rural and Environmental Biotechnology and Health Practice and Research, Department of Biological and Physical Sciences, University of Southern Queensland,² Queensland, Australia, and State Hygienic Laboratory, University of Iowa, Iowa City, Iowa³

Received 15 May 2000/Returned for modification 24 July 2000/Accepted 27 September 2000

No standardized PCR method is available for the laboratory diagnosis of the pertussis syndrome. Consensus recommendations for the use of PCR in the diagnosis of *Bordetella pertussis* infections have been proposed, and the aim of this study was to develop a method that fulfills all of these criteria. A rapid-cycle shared-primer PCR method with a microwell format and probe hybridization detection step (POR) was developed using novel oligonucleotides targeted to the outer membrane porin gene (*Bordetella* spp.). In specimens positive for *Bordetella* spp., *B. pertussis* was differentiated from *Bordetella parapertussis* and *Bordetella bronchiseptica* by hybridization with organism-specific oligonucleotide probes. An internal control was developed using overlap extension PCR and mouse β -actin DNA. The analytical specificity was 100%. The analytical sensitivity was comparable to that of nested IS481 and IS1001 PCR (~1 organism per reaction). The clinical sensitivity and specificity were ascertained using 705 specimens (from 705 patients). The results were compared to those of a nested-PCR method targeting the insertion sequences IS481 and IS1001. Fifty-one specimens were positive for *B. pertussis* by POR and IS481 PCR. Two specimens which fulfilled a clinical definition of pertussis were positive by POR and negative by IS481 PCR. A total of 652 specimens were negative by both methods. *B. parapertussis* was not detected in any specimens. PCR inhibition was detected in 21 out of 705 specimens (2.98%). Thus, a rapid (4 h, including specimen preparation) PCR method which fulfills all of the consensus recommendations was developed and validated for the detection of *B. pertussis*.

The diagnosis of *Bordetella pertussis* infections by nucleic acid amplification-based methods has, in general, been shown to be both highly sensitive and specific (13), yet an agreed-upon standardized method has not yet been adopted. The following consensus international recommendations have been published (11). (i) Sample processing should be kept to a minimum. (ii) Nasopharyngeal aspirates (NPA) are the preferred specimens. (iii) Differentiation between *B. pertussis* and *Bordetella parapertussis* is necessary. (iv) Carryover control (e.g., uracil-*N*-glycosylase) should be used to minimize contamination. (v) Appropriate controls (positive and negative) should be used. (vi) In detection systems, probes should be used for added specificity and sensitivity. (vii) Confirmation of questionable results with an alternative target is necessary. (viii) One hundred percent of culture-positive results should be nucleic acid amplification positive. (ix) Other commensals and pathogens should test negative. (x) Testing of samples from a healthy population assumed to be negative for the pathogen is required. (xi) Use of an internal control to check for inhibition of the reaction is required. However, as yet no method has been offered which fulfills all of these criteria. In a previous study (1), we developed a nested-PCR method utilizing the insertion sequences IS481 (*B. pertussis*) and IS1001 (*B. parapertussis*). Although the method had high analytical and clinical sensitivity and specificity and differentiated between *B. pertussis* and *B. parapertussis*, it was not useful for the routine diag-

nostic laboratory. Our efforts since then have been directed at developing a simpler method. In this study, we describe a rapid-cycle method which fulfills all of the consensus international recommendations and evaluate it using nested PCR utilizing both prospective specimens and specimens from a previous study (1). The results show that this method would be highly suited to serve as a standardized *B. pertussis* PCR method.

MATERIALS AND METHODS

Bacterial strains and genomic DNA. The bacterial strains and genomic DNA for sensitivity and specificity studies were the same as those described previously (1). Additional strains tested were *Bordetella avium* strain TC9 and *Bordetella hinzii* strain TC58, both kindly donated by Patrick Blackall from the Queensland Department of Primary Industries Animal Research Institute, Brisbane, Australia, and four *Bordetella holmesii* strains.

Patient specimens. Specimen collection, initial treatment, and storage were as previously described (1). A total of 705 specimens (from 705 patients) were tested. The specimen types were as follows: NPA, 608; throat (posterior pharyngeal) swabs, 82; and sputum, 15. Six hundred fifteen specimens were from patients with respiratory symptoms. Sixty-five NPA were from healthy asymptomatic adults, initially thought to be contacts in a pertussis outbreak which was later confirmed as an influenza A outbreak. Twenty-five throat swabs were collected from healthy adult volunteers. Samples were removed from the freezer (–20°C) and thawed at 37°C for 30 min. Twenty microliters was added to 80 μ l of sterile DNase- and RNase-free water, vortexed for 30 s, heated in a dry block heater for 20 min at 99.9°C, and then pulse centrifuged in a bench microcentrifuge (set at 11,300 \times g) for 30 s. Positive and negative controls were treated identically to specimens. Specimens in which inhibitory substances were detected (see the method described below) underwent a DNA extraction procedure using the High Pure PCR template kit (Boehringer Mannheim, Sydney, Australia) according to the manufacturer's instructions prior to retesting.

PCR methods. Nested IS481 PCR was performed as previously described (1). Pertussis toxin operon (PTO) PCR was also performed as described previously (D. J. Farrell, M. McKeon, G. Daggard, and T. K. S. Mukkur, Abstr. 39th

* Corresponding author. Present address: GR Micro Limited, 7-9 William Rd., London NW13ER, United Kingdom. Phone: 44 20 73887320. Fax: 44 20 73887324. E-mail: D.Farrell@grmicro.co.uk.

TABLE 2. Specificities of the individual probes in the POR assay for *Bordetella* spp.

Species	No. of strains positive for probe (target species)		
	DFPORP1 (<i>B. pertussis</i>)	DFPORP2 (<i>B. parapertussis</i> , <i>B. bronchiseptica</i>)	DFPORP3 (<i>B. pertussis</i> , <i>B. parapertussis</i> , <i>B. bronchiseptica</i>)
<i>B. pertussis</i> (21) ^a	21	0	21
<i>B. parapertussis</i> (6)	0	6	6
<i>B. bronchiseptica</i> (3)	0	3	3
<i>B. avium</i> (2)	0	0	0
<i>B. hinzii</i> (2)	0	0	0
<i>B. holmesii</i> (4) ^b	0	0	0
Non- <i>Bordetella</i> (30) (from reference 1)	0	0	0

^a Number of individual strains tested is given in parentheses.

^b Tested by the modified POR method in the Iowa laboratory.

washing and color development steps, the plates were read in a microwell plate reader at A_{450} .

Construction of an assay-specific internal control. In this study, mouse cDNA was prepared from BALB/c mouse spleens used for a different study. Mouse cDNA was used, as it was available; however, a simple DNA extract of mouse DNA would suffice. Hybrid primers (DFPORFMB and DFPORRMB) were designed using the sequence for mouse β -actin (GenBank accession no. X03672); the nucleotide sequences are listed in Table 1. A master mixture was prepared which had the following final concentrations: $1 \times$ PCR buffer (Life Technologies); 200 μ M (each) dATP, dCTP, dTTP, and dGTP (Pharmacia Biotech); 10 pmol of specific oligonucleotide per reaction mixture; platinum *Taq*, 2.0 U/reaction mixture; and 1.5 mM $MgCl_2$. Two microliters of mouse cDNA was added to 47.5 μ l of master mixture, and amplification was performed on a GeneAmp 9600 thermal cycler using the following parameters: 94°C for 2 min; 29 cycles of 94°C for 30 s, 55°C for 30 s, 72°C for 1 min; and 72°C for 5 min. Products were visualized on a 2% Tris-borate-EDTA agarose gel using ethidium bromide.

The resulting amplicon was of the same size as the amplicon for the *B. pertussis* assay and had the same sequences at either end but contained mouse DNA in the internal region. The internal control was then amplified by the diagnostic PCR method to produce a large amount of internal control. The internal control was then serially diluted and amplified in the standard *B. pertussis* assay, and the cutoff for positivity was determined. One dilution higher than the cutoff concentration was the concentration used in the master mixture. If PCR-inhibitory substances were present in the specimen, they would be detected by failure to amplify the internal control, which was detected in a separate microwell using an oligonucleotide probe (DFMBAP) specific for mouse β -actin DNA.

RESULTS

Analytical specificity was 100% using the strains previously described (1), *B. avium* strain TC9, *B. hinzii* strain TC58, and four *B. holmesii* strains (Table 2). No positive results were obtained for 65 NPA or 25 throat swabs collected from 90 asymptomatic healthy adults. The analytical sensitivity of the assay was between 1 and 10 CFU per reaction mixture. Table 3 shows that analytical sensitivity was not adversely affected for *B. pertussis* or *B. parapertussis* if either organism was present in minimal numbers and the other in large numbers.

In the comparison between POR and IS481 nested PCR for 705 clinical specimens, 51 specimens were positive and 652 specimens were negative for both tests. No tests were positive for IS481 and negative for POR; however, two POR-positive and IS481-negative specimens were obtained. The two specimens that produced discrepant results came from two patients who fulfilled a clinical definition of pertussis (14) and were also positive by PTO PCR. PCR inhibition was detected in 21 of 705 specimens (2.98%). Inhibitors were detected more frequently in extremely mucoid specimens, of which 8 were sputum and 13 were NPA. After DNA extraction, no inhibitors were detected for any of the 21 specimens. One of the inhib-

itory specimens was positive for *B. pertussis* after DNA extraction.

DISCUSSION

Disadvantages of the nested IS481-IS1001 method arise from the inherent problems of nested-PCR assays, in particular, the increased risk of carryover of amplified DNA. In addition, the carryover cannot be minimized by the use of the uracil-*N*-glycosylase carryover prevention system. Because of these problems, this method should only be performed in laboratories with staff experienced in nested-PCR techniques, and hence it is not suitable for routine laboratory use. The next step was to develop a simpler assay, using the IS481-IS1001 assay as a "gold standard" to evaluate the analytical and clinical parameters.

The two main candidates considered as appropriate targets were the repetitive insertion sequence IS481 and the regulatory region of the pertussis toxin gene. IS481 is present in all strains of *B. pertussis* tested so far at a rate of 50 to 100 copies per chromosome (1, 9, 10). IS481 would seem like an appropriate target due to its high copy number, yet many have questioned its use. Grimprel et al. (3) argued that the copy number might vary between strains and that insertion sequences may also be present in other *Bordetella* species. To assess the specificity of the insertion sequence, Glare et al. (2) performed hybridization studies with 45 different species of bacteria (from 24 genera) using a *Clal* fragment as a probe. Strong hybridization occurred for *B. pertussis*, weak hybridization was observed for *B. bronchiseptica* (after prolonged autoradiography), and no hybridization was observed for any other species. In all studies except that of Glare et al. (2), the specificity of IS481-based PCR has been 100% when tested against a multitude of non-pertussis bordetellae, and a high correlation has been observed with clinical pertussis and culture. On the other hand, Glare et al. (2) reported a 12-bp difference compared to the sequence published by McLafferty et al. (9), suggesting that the insertion sequence is not absolutely conserved. Also, a recent study (8) showed that an IS481 PCR cross-reacted strongly with six strains of *B. holmesii*. Unfortunately, the carriage rate and clinical significance of *B. holmesii* are unknown, confounding assays that target IS481.

In the first attempt to develop a simpler assay, the pertussis toxin operon was used as a target, and the PTO assay performed well, both analytically and clinically, against nested IS481-IS1001 (Farrell et al., Abstr. 39th Intersci. Conf. Antimicrob. Agents Chemother.). Although *ptx* is present as a single-copy gene, a comparison of PTO and IS481 as targets showed equivalent sensitivities (4). On the other hand, identical PCR protocols were used for both IS481 and PTO, yet the

TABLE 3. Absorbances for reactions containing different amounts of *B. pertussis* (strain Tohama I) and *B. parapertussis* (strain ATCC 15237)

<i>B. pertussis</i>	No. of bacteria ^a	$A_{450/620}$ ^b			
		<i>B. parapertussis</i>	DFPOR1	DFPOR2	DFPOR3
1	10^7		0.03	2.2	2.25
10	10^7		1.2	2.3	2.1
100	10^5		2.25	2.35	2.4
10^3	10^3		2.3	2.15	2.25
10^5	100		2.15	1.95	2.1
10^7	10		2.4	1.6	2.35
10^7	1		2.25	0.25	2.05

^a CFU per reaction mixture.

^b Average of three readings.

published melting temperatures (T_m) of the primers differed by 14 and 16°C between the two methods (4). One could argue that with better optimization the IS481 PCR might have been a great deal more sensitive than PTO PCR. There are arguments for not using PTO as a target. (i) The pertussis toxin gene is also present but not expressed in *B. parapertussis* and *B. bronchiseptica* due to multiple mutations located in the *ptx* promoter region (6). On the other hand, an immunoglobulin G response to pertussis toxin in patients with parapertussis infections has been described, indicating that expression of the toxin genes may occur (5). (ii) Mutations in the pertussis toxin S1 subunit have recently been described; Mooi et al. (12) provide evidence that contemporary strains of *B. pertussis* differ in the genetic sequence of the S1 subunit, possibly through vaccine-driven evolution. For diagnostic methods, it may be wise not to use genes that code for antigens present in commercial vaccines. In addition, polymorphisms in the pertussis toxin promoter region have also been described (15).

The region upstream of the outer membrane porin gene has been used as a target for diagnostic PCR in only one previous study (5). The assay described in that study had a sensitivity of 79% against culture when gel detection was used and a sensitivity of 95% when DIG immunoblotting was used. In the present study, new primers and probes were designed using the outer membrane porin sequence data described by Li et al. (5). A rapid-cycle protocol was designed; a carryover prevention system (uracil-*N*-glycosylase) was included; probe detection allowed differentiation among *Bordetella* spp.; an assay-specific internal control was developed, allowing specimen preparation to be minimal; and the assay was evaluated against a gold standard method targeting a different gene. This method is the first to fulfill the international consensus requirements for the use of PCR in the detection of *B. pertussis*, as proposed by Meade and Bollen (11) and outlined above. The assay does not distinguish between *B. parapertussis* and *B. bronchiseptica*, but this should only cause concern in interpreting results from specimens obtained from immunocompromised patients, as *B. bronchiseptica* is only rarely part of the normal flora of humans. Nevertheless, interpretation of a positive *B. parapertussis*-*B. bronchiseptica* POR result should be considered with caution. The POR PCR assay allows a rapid, accurate, standardized, and simplified approach to the laboratory diagnosis of pertussis.

ACKNOWLEDGMENTS

This study was funded in part by the Private Practice Trust Fund, Toowoomba Health Region (D.J.F.), and the Project Team Research Project Grant of the University of Southern Queensland (T.K.S.M.).

The support of the pediatricians and general practitioners from the Darling Downs region is gratefully acknowledged. Specimens were from the Queensland Health Pathology Service Gold Coast (David Alfredson), Townsville and Mount Isa Laboratories, and Sullivan and Nicolaides and Partners Laboratories, Brisbane and Lismore (Lynne Wright). Strains of bacteria were from J. Faoagali, M. Nolan, J. Bates, R. Reed, R. McDougall, C. Coulter, Pat Blackall, and Alison Weiss.

REFERENCES

- Farrell, D. J., G. Daggard, and T. K. S. Mukkur. 1999. Nested duplex PCR to detect *Bordetella pertussis* and *Bordetella parapertussis* and its application in diagnosis of pertussis in nonmetropolitan South East Queensland, Australia. *J. Clin. Microbiol.* **37**:606–610.
- Glare, E. M., J. C. Paton, R. R. Premier, A. J. Lawrence, and I. T. Nisbet. 1990. Analysis of a repetitive DNA sequence from *Bordetella pertussis* and its application to the diagnosis of pertussis using the polymerase chain reaction. *J. Clin. Microbiol.* **28**:1982–1987.
- Grimprel, E., P. Bégué, I. Anjak, F. Betsou, and N. Guiso. 1993. Comparison of polymerase chain reaction, culture, and Western immunoblot serology for diagnosis of *Bordetella pertussis* infection. *J. Clin. Microbiol.* **31**:2745–2750.
- Houard, S., C. Hackel, A. Herzog, and A. Bollen. 1989. Specific identification of *Bordetella pertussis* by the polymerase chain reaction. *Res. Microbiol.* **140**:477–487.
- Li, Z., D. L. Jansen, T. M. Finn, S. A. Halperin, A. Kasina, S. P. O'Connor, T. Aoyama, C. R. Manclark, and M. J. Brennan. 1994. Identification of *Bordetella pertussis* infection by shared-primer PCR. *J. Clin. Microbiol.* **32**:783–789.
- Locht, C., and J. M. Keith. 1986. Pertussis toxin gene: nucleotide sequence and genetic organization. *Science* **232**:1258–1264.
- Loeffelholz, M. J., C. J. Thompson, K. S. Long, and M. J. R. Gilchrist. 1999. Comparison of PCR, culture, and direct fluorescent-antibody testing for detection of *Bordetella pertussis*. *J. Clin. Microbiol.* **37**:2872–2876.
- Loeffelholz, M. J., C. J. Thompson, K. S. Long, and M. J. R. Gilchrist. 2000. Detection of *Bordetella holmesii* using *Bordetella pertussis* IS481 PCR assay. *J. Clin. Microbiol.* **38**:467.
- McLafferty, M. A., D. R. Harcuss, and E. L. Hewlett. 1988. Nucleotide sequence and characterization of a repetitive element from the genome of *Bordetella pertussis* with characteristics of an insertion sequence. *J. Gen. Microbiol.* **134**:2297–2306.
- McPheat, W. L., and T. McNally. 1987. Isolation of a repeated DNA sequence from *Bordetella pertussis*. *J. Gen. Microbiol.* **133**:323–330.
- Meade, B. D., and A. Bollen. 1994. Recommendations for the use of the polymerase chain reaction in the diagnosis of *Bordetella pertussis* infections. *J. Med. Microbiol.* **41**:51–55.
- Mooi, F. R., H. van Oirschot, K. Heuvelman, H. G. J. van der Heide, W. Gaastra, and R. J. L. Willems. 1998. Polymorphism in the *Bordetella pertussis* virulence factors P.69/pertactin and pertussis toxin in the Netherlands: temporal trends and evidence for vaccine-driven evolution. *Infect. Immun.* **66**:670–675.
- Müller, F. C., J. E. Hoppe, and C. H. Wirsing von König. 1997. Laboratory diagnosis of pertussis: state of the art in 1997. *J. Clin. Microbiol.* **35**:2435–2443.
- National Health and Medical Research Council. 1994. Surveillance case definitions. National Health and Medical Research Council, Canberra, Australia.
- Nygren, M., E. Reizenstein, M. Ronaghi, and J. Lundeberg. 2000. Polymorphism in the pertussis toxin promoter region affecting the DNA-based diagnosis of *Bordetella* infection. *J. Clin. Microbiol.* **38**:55–60.