

Effects of HIF-1 α on ERR α / γ protein expression in mouse skeletal muscle

Ying Zhang, Weixiu Ji, Lianfeng Zhang, Sixue Liu, Gang Liu, Jianxiong Wang

ABSTRACT

Aims: HIF-1 α plays an important role in the adaptive responses to hypoxia. The ERR α and γ are crucial regulators of energy metabolism in skeletal muscle. The aim of the present study was to generate the inducible HIF-1 α transgenic mice and examine the effects of different HIF-1 α protein expression levels on ERR α / γ in mouse skeletal muscle. **Methods:** We generated the HIF-1 α high-expression transgenic mice (HT) and HIF-1 α low-expression transgenic mice (LT), and then compared the expressions of ERR α / γ and its target genes in skeletal muscles of three kinds of mice: HT, LT, and non-transgenic mice (NT). **Results:** The results showed that (1) the double positive mice from the founder of 3# and 68# showed an obvious expression of HIF-1 α induced by tamoxifen and both of them were maintained to the further research as HT and LT mice, respectively; and (2) the nucleoprotein expressions of ERR α / γ and the mRNA levels of

the ERR α / γ target genes: MCAD, PPAR α , NRF1 and PDK4 were higher in the LT mice than the values in the NT, but only the mRNA levels of MCAD and PPAR α were significantly higher. The HT mice showed significantly lower ERR α protein content than that of the NT mice. **Conclusion:** Our study was the first to report the generation of the inducible HIF-1 α transgenic mice and effects of HIF-1 α on ERR α / γ protein expression in mouse skeletal muscle in vivo. These data demonstrate that the low HIF-1 α protein expression may associate with an up-regulation of ERR α / γ and their target genes in skeletal muscles, while the aggravated HIF-1 α protein expression would reduce the effects.

Keywords: ERR α / γ , HIF-1 α , Skeletal muscle, Target genes, Transgenic mouse,

How to cite this article

Zhang Y, Ji W, Zhang L, Liu S, Liu G, Wang J. Effects of HIF-1 α on ERR α / γ protein expression in mouse skeletal muscle. Edorium J Biomed Technol 2015;1:4–10.

Article ID: 100002B05YZ2015

doi:10.5348/B05-2015-2-OA-2

Ying Zhang¹, Weixiu Ji¹, Lianfeng Zhang², Sixue Liu¹, Gang Liu³, Jianxiong Wang⁴

Affiliations: ¹Institute of Sports Science, Beijing Sport University, China; ²Key Laboratory of Human Disease Comparative Medicine, Ministry of Health, Institute of Laboratory Animal Science, Chinese Academy of Medical Sciences & Comparative Medical Center, Peking Union Medical College, China; ³Teaching Laboratory Centre, Beijing Sport University, China; ⁴School of Health and Wellbeing, Faculty of Health, Engineering, and Sciences, University of Southern Queensland, Australia.

Corresponding Author: Professor Ying Zhang, Institute of Sport Science, Beijing Sport University, Beijing 100084 China; Ph: +86 10 62989584; Fax: +86 10 62989444; Email: zhyi9256@126.com

Received: 24 September 2015

Accepted: 28 October 2015

Published: 03 February 2016

INTRODUCTION

The hypoxia inducible factor (HIF)-1 plays a key role to mediate the cellular responses to hypoxia. Its target genes control many cellular signal transductions, such as erythropoietin which induces red blood cell

production, vascular endothelial growth factor (VEGF) which promotes angiogenesis, and glucose transporter 1 (GLUT1) which increases the efficiency of the glucose uptake [1, 2]. Previous studies have examined the role of HIF-1 α in skeletal muscle. A research group found that the HIF-1 protein and mRNA levels are varied in different fiber types of rodent skeletal muscles [3]. Another study demonstrated that acute exercise can elevate the HIF-1 protein level in human muscle [4]. Deletion of HIF-1 in mouse skeletal muscle might lead to impaired glycolytic flux and decreased exercise endurance [5]. These studies focused on the effects of HIF-1 on the tissue or organ level. Skeletal muscle function and metabolism are controlled by many signaling pathways and molecules. At the molecular level, there are few studies about the relationship between HIF-1 α and other signaling pathways or molecules which control energy metabolism in skeletal muscle.

Estrogen-related receptors (ERRs) are crucial regulators of energy metabolism [6]. There are three members in the ERR family: ERR α , ERR β and ERR γ . They are active nuclear receptors that contain high levels of the DNA sequence identified to that of estrogen receptors [7]. ERR α and γ isoforms are ubiquitously expressed at high levels in the heart, skeletal muscle and brown adipose tissue; while ERR β expression is restricted to the brain, kidney and heart. ERRs control many downstream target genes which are important to glycolytic metabolism, fatty acid oxidation, mitochondrial oxidative phosphorylation, and mitochondrial biosynthesis. The main downstream target genes include pyruvate dehydrogenase kinase 4 (PDK4) [8], medium-chain acyl coenzyme A dehydrogenase (MCAD) [9], peroxisome proliferator-activated receptor α (PPAR α) [10], and nuclear respiratory factors (NRFs) [11].

There are a few reports about the relationship between HIF-1 and ERRs in cell lines in literature. A previous study has shown that ERR γ mRNA and protein levels were increased by a hypoxic treatment in the hepatoma cell lines and this change in ERR γ was associated with the increased HIF-1 α and β expression [12]. The knockdown of endogenous HIF-1 α reduced the hypoxia-mediated induction of ERR γ . In addition, hypoxia also increased the promoter activity and mRNA level of PDK4 in the HepG2 cells [12]. Another study suggested that ERRs physically interact with HIF and stimulate HIF-induced transcription. Transcriptional activation of hypoxic genes in cells cultured under hypoxia was largely blocked by suppression of ERRs through the treatment with diethylstilbestrol, an ERR inhibitor [13]. These results have suggested a possible relationship between HIF-1 α and ERRs. However, these observations are restricted to the cell culture studies. There are few researches of the relationship between HIF1 α and ERR α/γ in vivo. The purpose of this study was to generate the inducible HIF-1 α transgenic mice and evaluate the effects of HIF-1 α expression on the ERR α/γ protein and the mRNA

expression levels of the ERR α/γ 's target genes in skeletal muscle of the mouse with low- or high-expression in HIF- α protein.

MATERIALS AND METHODS

The generation of the inducible HIF-1 α transgenic mice

The eGFP gene was labeled with two LoxP sites in the 5' and 3' end and fused with a full length mouse HIF-1 α cDNA (Genbank accession No.15251) (Figure 1). The transgenic plasmid (pBactin2(+)-GFP-Loxp-mHIF-1 α) was constructed by inserted the DNA fragment containing the eGFP and HIF-1 α cDNA under the chicken β -actin promoter, that the chicken β -actin promoter enhanced the eGFP expression directly and could enhance the HIF-1 α expression after the eGFP to be removed by Cre. The inducible expression of HIF-1 α mRNA was tested in the 293T cells (Figure 2). The 293T cells were transfected with pcDNA3.1(-)-CRE, pBactin2(+)-GFP-Loxp-mHIF-1 α , and pcDNA3.1(-)-CRE + pBactin2(+)-GFP-Loxp-mHIF-1 α , respectively. The total RNA was isolated and mRNA levels of the HIF-1 α were detected by RT-PCR. The construct was microinjected into male pronuclei of fertilized mouse oocytes and implanted into pseudo-pregnant females to generate the transgenic mouse lines [14]. The Genotyping of GFP-Loxp-mHIF-1 α transgenic mice was verified by PCR using primers 5'TGGCAATGTCTCCTTTACCTTC and 5'GCAGTGGTCGTTTCTTGAGG to amplify the desired 654bp fragment of the mHIF-1 α (Figure 3). Totally, 12 founder mice were found and they were crossed with a pUbc-Cre-ert2 mice (from JAX, No.008085), which could expressed the Cre-ert2 systematically in an inducible way with tamoxifen (Figure 4). The double positive mice were generated from 2#, 3#, 7# and 68# founders and




Figure 1: A LoxP labeled eGFP between the chicken β -action promoter and the HIF-1 α cDNA.




Figure 2: The HIF-1α expression in 293T cells.




Figure 3: The Genotyping of GFP-Loxp-mHIF-1α transgenic mice was verified by PCR.




Figure 4: There were four double positive F1 mice from the 12 founder crossed with the pUBC-Cre-ert2 mice respectively.

these mice were induced with tamoxifen in a dose of 100 μg/d by oral administration for 10 days to get HIF-1α low-expression (LT) and HIF-1α high-expression (HT) transgenic mice. The double positive mice were maintained on a C57BL/6J genetic background and all the mice were bred in an AAALAC-accredited facility.

The present study protocol was approved by the Animal Care and Use Committee of Beijing Sport University. Non-transgenic mice (NT) (n=10), LT mice (n=10), and HT mice (n=10) were used to evaluate the effects of HIF-1α expression on the ERRα/γ protein and the mRNA expression levels of the ERRα/γ's target genes in skeletal muscle of the mouse. All mice with a mean body weight of 20±2 g were housed with controlled room temperature (20–25°C), 12:12-h light-dark cycle, and free access to food and water. After allowing acclimatization to their housing for three days, all mice were euthanized and their skeletal muscles from the legs were excised, cleaned of blood and connective tissue, quick-frozen with aluminum tongs, pre-cooled in liquid nitrogen, and stored at -80°C.

Western blot analyses

For Western blot analyses of HIF-1α, ERRα and ERRγ protein levels, homogenates were made from the skeletal muscle samples. Nucleoproteins were isolated from 100 mg of muscle using NE-PER nuclear extraction reagents (87792, Pierce, Rockford, IL, USA). Protein concentration was measured using the BCA protein assay kit (23225, Pierce, Rockford, IL, USA). Nucleoproteins (20 μg) were separated on NuPAGE®Novex®Bis-Tris Gels (separating gel 4–12%) (Life Technologies, USA) by electrophoresis, and the fractionated proteins were transferred to a nitrocellulose membrane by iBlot Gel Transfer System (Invitrogen, USA). The blots were probed with HIF-1α antibody (Cell signaling, 3434), ERRα antibody (abcam, 76228), ERRγ antibody (abcam, 128930), and the same blots were stripped and reprobed with GAPDH (Santa Cruz, sc-48166)/β-actin (Santa Cruz, sc-47778) or histone-1 (H1) antibody (Santa Cruz, sc-8030). The individual values were originally expressed as a ratio of a standard (GAPDH/β-actin or H1 content) and then expressed as a fold change of the NT group value.

Real-time PCR

Total RNA was isolated from 50 mg of crushed muscle tissue using the TRI reagents according to the manufacturer's instructions. Reverse transcription (RT) of total RNA to cDNA was performed using a Reverse Transcription Kit (Toyobo, Japan). RT-PCR was performed in a ABI 7500 Real-Time PCR System (USA) using 10 ul of SYBR Green Real time PCR Master Mix (Toyobo, Japan), 2 ul of cDNA template, 6 ul of ddH₂O and 2ul of respective Qiagen primer sets (Germany) for PDK4 (QIAGEN, QT00157248), MCAD (QIAGEN, QT00111244), PPARα (QIAGEN, QT00137984), NRF1 (QIAGEN, QT01051820) and HIF-1α (QIAGEN, QT01039542). Copy numbers of cDNA targets were quantified using Ct values, and the mRNA expression levels for all samples were normalized to the level of the housekeeping gene 18S rRNA (QIAGEN, QT010036875). The difference in expression between control and experimental was calculated as $2^{-\Delta\Delta CT}$, as described previously [15].

Statistical analysis

All data were presented as means ± SE. Statistical analysis was conducted via one-way ANOVA, and a Tukey post hoc test was used when significance was found. Differences were considered statistically significant at $p < 0.05$.

RESULTS

The generation of the inducible HIF-1 α transgenic mice

The generation of the inducible HIF-1 α transgenic mice was shown in Figure 5. The double positive mice from the founder of 3# and 68# showed an obvious expression of HIF-1 α induced by tamoxifen and both of them were maintained to the further research as LT and HT transgenic mice, respectively.

HIF-1 α and nucleoprotein ERR α expressions in skeletal muscle

Skeletal muscles of NT, LT and HT mice were harvested and their expression of HIF-1 α and ERR α were analyzed by Western blot. LT and HT mice showed higher HIF-1 α signals than that of the NT mice and HT mice showed higher HIF-1 α signal than that of the LT mice when using the HIF-1 α antibody in muscle (Figure 6A). The nucleoprotein expression of ERR α in LT mouse skeletal muscle was increased, but did not reach significant level; while the nucleoprotein expression of ERR α in HT mouse skeletal muscle was decreased significantly, compared with that of the NT muscles (Figure 6B).

The nucleoprotein expression of ERR γ in skeletal muscle

There was a trend that the nucleoprotein expression of ERR γ was higher in the LT group and was lower in the HT group, compared to that of the NT group, but the differences were not significant between the groups (Figure 7). The ERR γ nucleoprotein level in LT and HT groups had a similar change pattern of the ERR α in Figure 6.

The mRNA expression of MCAD, NRF1, PPAR α and PDK4 in skeletal muscle

The levels of MCAD, NRF1, PPAR α and PDK4 mRNA expressions were measured using RT-PCR and normalized with 18S rRNA in the three groups (Figure 4). The mRNA expressions of MCAD and PPAR α in the LT mice were significantly increased, compared with the value in the NT. The mRNA expressions of NRF1 and PDK4 showed increases but the trends were not statistically significant. The HT mice did not show significant difference in the genes' mRNA expressions to those of the LT and NT mice (Figure 8).




Figure 5: The inducible expression of HIF-1 α in tissue of muscle was compared with Western blot.




Figure 6: HIF-1 α protein content (A) and ERR α nucleoprotein content, (B) in NT, LT and HT mice muscles.

*significantly different from the corresponding NT mice values at $p < 0.05$.




Figure 7: Western blot analysis for ERR γ in mouse skeletal muscle.




Figure 8: The levels of MCAD, NRF1, PPAR α and PDK4 mRNA. * significantly different from the corresponding NT mice values at $p < 0.05$.

DISCUSSION

The importance of HIF-1 signaling in metabolic regulation has spurred much interest in recent years [12, 16]. Previous studies have been mainly focused on the metabolic role of HIF in

- (i) stimulating the expression of genes that encodes glucose transporters and glycolytic enzymes to accelerate glucose uptake and the anaerobic glycolytic ATP production [17],
- (ii) down-regulating mitochondrial oxygen consumption by directly or indirectly inducing pyruvate dehydrogenase (PDK), which inhibits the mitochondrial pyruvate dehydrogenase complex (PDC) from converting pyruvate into acetyl-CoA [18,19], and
- (iii) regulating the differential expression of cytochrome c oxidase subunit 4 isoforms to optimize the efficiency of respiration at different O₂ tensions [20].

There are few studies referred to the relationship of HIF and ERR. Studying the effects of HIF-1 α on ERR α/γ would improve our understanding about the metabolic regulation under hypoxic condition in skeletal muscle. Therefore, in the present study for the first time, we generated the inducible HIF-1 α transgenic mice and reported the impact of HIF-1 α on ERR α/γ protein expression in mouse skeletal-muscle through the new generated HIF-1 α transgenic mice.

There are no conclusive results of the effects of HIF-1 on ERRs in hypoxia. Some studies have reported that hypoxia stimulates the expression of ERR α through peroxisome proliferator-activated receptorcoactivator-1 α (PGC-1 α), but not HIF-1. [21, 22]. While another study showed that HIF-1 α had an effect on ERRs in hypoxic cell culture in vitro [12]. This statement has been supported by the finding of two putative binding sites of HIF-1 α on the ERR γ promoter in a cell culture

experiment. The promoter activity in response to hypoxia was abolished in these site-mutated constructs [12]. Furthermore, the evidence of ERR γ regulation of HIF-1 α under hypoxic condition was confirmed by the ChIP assays in the hypoxia treated HepG2 [12]. In the present study, we observed that the nucleoprotein expressions of ERR α/γ and the mRNA levels of the ERR α/γ target genes: MCAD, PPAR α , NRF1 and PDK4 were higher in the LT mice compared to the values in the NT mice, although only the mRNA levels of MCAD and PPAR α were increased significantly. The results suggest that HIF and ERR may act in a common pathway and form a regulatory complex. Interestingly, we also found that the ERR α/γ protein contents were decreased in the HT mice and the reduced ERR α was significantly different from the corresponding NT value. HIF-1 α is able to induce apoptosis during severe or prolonged (>24 hours) hypoxia, although it has been generally accepted as a pro-survival factor [13, 23]. Thus, the molecular mechanism causing different ERR α/γ protein levels, transcriptional regulation or output of ERR α/γ for downstream targets requires further clarification.

There are limitations in our study, we had no single transgenic mice of GFP-Loxp-mHIF-1 α and Ubc-Cre-ert2 as controls. More controls would provide more clear results to the research. In addition, we did not measure the transcriptional activity of the ERR α/γ genes which are regulated by HIF-1 α , as well as the ERR α/γ mRNA level and the transcriptional activity of PDK4, MCAD, PPAR α and NRF1 genes which are regulated by ERR α/γ . Though the present results have shown the effects of HIF-1 α on the expressions of ERR α/γ and their target gene, future studies should measure these directly. Moreover, in the present study, we only investigated the gene and protein expressions but did not do any histomorphological and functional measurements. These histomorphological and functional measurements would be desirable and add more comprehensive insights into the scope of the study.

CONCLUSION

Our study was the first to report the generation of the inducible HIF-1 α transgenic mice and effects of HIF-1 α on ERR α/γ protein expression in mouse skeletal muscle in vivo. We found that the nucleoprotein expressions of ERR α/γ and the mRNA levels of the ERR α/γ target genes were higher in the LT mice; while the ERR α/γ protein contents were decreased in HT mice, compared to the values in the NT mice respectively. These data reveal a relationship that HIF-1 α affects ERR α/γ and their target genes in mouse skeletal muscle.

Acknowledgements

This study was funded by grants from the National Natural Science Foundation of China (31471134; 31171140) and

Program of General Administration of Sport of China (2014B010).

Author Contributions

Ying Zhang – Substantial contributions to conception and design, Acquisition of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Weixiu Ji – Substantial contributions to conception and design, Acquisition of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Lianfeng Zhang – Substantial contributions to conception and design, Acquisition of data, Drafting the article, Final approval of the version to be published

Sixue Liu – Substantial contributions to conception and design, Acquisition of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Gang Liu – Substantial contributions to conception and design, Acquisition of data, Drafting the article, Final approval of the version to be published

Jianxiong Wang – Substantial contributions to conception and design, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Guarantor

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.

Copyright

© 2015 Ying Zhang et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

REFERENCES

1. Nakayama K. Cellular signal transduction of the hypoxia response. *J Biochem* 2009 Dec;146(6):757–65.
2. Ke Q, Costa M. Hypoxia-inducible factor-1 (HIF-1). *Mol Pharmacol* 2006 Nov;70(5):1469–80.
3. Pisani DF, Dechesne CA. Skeletal muscle HIF-1 α expression is dependent on muscle fiber type. *J Gen Physiol* 2005 Aug;126(2):173–8.
4. Ameln H, Gustafsson T, Sundberg CJ, et al. Physiological activation of hypoxia inducible factor-1 in human skeletal muscle. *FASEB J* 2005 Jun;19(8):1009–11.
5. Mason SD, Howlett RA, Kim MJ, et al. Loss of skeletal muscle HIF-1 α results in altered exercise endurance. *PLoS Biol* 2004 Oct;2(10):e288.
6. Ariazi EA, Jordan VC. Estrogen-related receptors as emerging targets in cancer and metabolic disorders. *Curr Top Med Chem* 2006;6(3):203–15.
7. Giguère V. Transcriptional control of energy homeostasis by the estrogen-related receptors. *Endocr Rev* 2008 Oct;29(6):677–96.
8. Zhang Y, Ma K, Sadana P, et al. Estrogen-related receptors stimulate pyruvate dehydrogenase kinase isoform 4 gene expression. *J Biol Chem* 2006 Dec 29;281(52):39897–906.
9. Sladek R, Bader JA, Giguère V. The orphan nuclear receptor estrogen-related receptor alpha is a transcriptional regulator of the human medium-chain acyl coenzyme A dehydrogenase gene. *Mol Cell Biol* 1997 Sep;17(9):5400–9.
10. Huss JM, Torra IP, Staels B, Giguère V, Kelly DP. Estrogen-related receptor alpha directs peroxisome proliferator-activated receptor alpha signaling in the transcriptional control of energy metabolism in cardiac and skeletal muscle. *Mol Cell Biol* 2004 Oct;24(20):9079–91.
11. Rangwala SM, Li X, Lindsley L, et al. Estrogen-related receptor alpha is essential for the expression of antioxidant protection genes and mitochondrial function. *Biochem Biophys Res Commun* 2007 May 25;357(1):231–6.
12. Lee JH, Kim EJ, Kim DK, et al. Hypoxia induces PDK4 gene expression through induction of the orphan nuclear receptor ERR β . *PLoS One* 2012;7(9):e46324.
13. Ao A, Wang H, Kamarajugadda S, Lu J. Involvement of estrogen-related receptors in transcriptional response to hypoxia and growth of solid tumors. *Proc Natl Acad Sci U S A* 2008 Jun 3;105(22):7821–6.
14. Gordon JW, Ruddle FH. Integration and stable germ line transmission of genes injected into mouse pronuclei. *Science* 1981 Dec 11;214(4526):1244–6.
15. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001 Dec;25(4):402–8.
16. Formenti F, Constantin-Teodosiu D, Emmanuel Y, et al. Regulation of human metabolism by hypoxia-inducible factor. *Proc Natl Acad Sci U S A* 2010 Jul 13;107(28):12722–7.
17. Firth JD, Ebert BL, Pugh CW, Ratcliffe PJ. Oxygen-regulated control elements in the phosphoglycerate kinase 1 and lactate dehydrogenase A genes: similarities with the erythropoietin 3' enhancer. *Proc Natl Acad Sci U S A* 1994 Jul 5;91(14):6496–500.
18. Papandreou I, Cairns RA, Fontana L, Lim AL, Denko NC. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* 2006 Mar;3(3):187–97.
19. Kim JW, Tchernyshyov I, Semenza GL, Dang CV. HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab* 2006 Mar;3(3):177–85.
20. Fukuda R, Zhang H, Kim JW, Shimoda L, Dang CV, Semenza GL. HIF-1 regulates cytochrome oxidase subunits to optimize efficiency of respiration in hypoxic cells. *Cell* 2007 Apr 6;129(1):111–22.
21. Zhang K, Lu J, Mori T, et al. Baicalin increases VEGF expression and angiogenesis by activating the

- ERR α /PGC-1 α pathway. *Cardiovasc Res* 2011 Feb 1;89(2):426–35.
22. Arany Z, Foo SY, Ma Y, et al. HIF-independent regulation of VEGF and angiogenesis by the transcriptional coactivator PGC-1 α . *Nature* 2008 Feb 21;451(7181):1008–12.
23. Greijer AE, van der Wall E. The role of hypoxia inducible factor 1 (HIF-1) in hypoxia induced apoptosis. *J Clin Pathol* 2004 Oct;57(10):1009–14.

Access full text article on
other devices



Access PDF of article on
other devices

