

1 **Anaerobic toxicity assay of plasticisers**  
2

3  
4  
5  
6 ANTOINE P. TRZCINSKI, NKECHI OFOEGBU and DAVID C. STUCKEY\*  
7

8  
9 *Department of Chemical Engineering, Imperial College of Science and Technology*  
10 *and Medicine, Prince Consort Road, London SW7 2BY, UK*  
11

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32 \_\_\_\_\_  
33 \*Address correspondence to Professor David C. Stuckey, Department of Chemical  
34 Engineering, Imperial College of Science and Technology and Medicine, Prince  
35 Consort Road, London SW7 2BY, UK; Tel.: +44 (0)20 7594 5591; Fax: +44 (0)20  
36 7594 5629  
37 E-mail: [d.stuckey@ic.ac.uk](mailto:d.stuckey@ic.ac.uk) or [a.trzcinski05@ic.ac.uk](mailto:a.trzcinski05@ic.ac.uk)

38

39

40

41

42

43 **ABSTRACT**

44

45 Plasticisers are commonly found in landfill leachate and accumulate in the  
46 environment. Some of them are known as disruptive endocrine compound. This  
47 manuscript assessed the toxicity of three common plasticisers, including Bis(2-  
48 Ethylhexyl)phthalate (DEHP), o-hydroxybiphenyl (HBP) and 2,6-di-tert-butyl-4-  
49 (dimethylaminomethyl) phenol (MAMP) on the methanogens during the anaerobic  
50 process. It was found that DEHP and MAMP did not impede methanogenesis up to  
51 200 mg/L, but no additional methane could be obtained from their degradation. In  
52 contrast, HBP severely inhibited methanogens at 200 mg/L, but after acclimatisation  
53 it could be metabolised resulting in a 25 % increase in methane production compared  
54 to the control.

55

56 **Keywords:** Anaerobic toxicity assay; plasticiser; phthalate.

57

58

59 **INTRODUCTION**

60

61

62 Landfill is one the most widely employed methods for the disposal of municipal solid  
63 waste around the world. Landfill leachate can be defined as the rain water percolating  
64 through landfills and containing harmful substances for the environment. Plastics can  
65 accumulate in the landfill because of their non-biodegradability, and components  
66 from the plastic can leach out and be found in landfill leachate. These toxic  
67 components are used during the manufacturing of plastics to improve their  
68 processability and are called plasticisers. These plasticisers can also be found on  
69 industrial sites and some are known to accumulate in the environment due to human  
70 activities.

71 Common plasticisers are o-hydroxybiphenyl (HBP) and Bis(2-Ethylhexyl)phthalate  
72 (DEHP). HBP consists of two linked benzene rings and a phenolic hydroxyl group,  
73 and is on the list of chemicals recognized as carcinogens by the state of California. <sup>[1]</sup>  
74 It was found at a concentration of 23 µg/L in the effluent of our anaerobic membrane  
75 bioreactor treating simulated municipal solid waste, <sup>[2]</sup> but some authors have reported  
76 a concentration of 2 µg/L in actual landfill leachate. <sup>[3]</sup> DEHP is the most important  
77 phthalate, is produced on a massive scale due to its good plasticizing properties, and is  
78 known to be a disruptive endocrine compound and has carcinogenic and mutagenic  
79 effects. <sup>[4-5]</sup> DEHP was reported to be barely biodegradable under anaerobic  
80 conditions, <sup>[6]</sup> while other authors <sup>[7]</sup> found that it could not be removed by aeration,  
81 coagulation/sedimentation or biological treatment. Gavala et al. <sup>[5]</sup> showed that  
82 degradation of DEHP occurred in a digester treating primary sludge, but accumulation  
83 of high levels of DEHP (more than 60 mg/L) had a negative effect on DEHP removal  
84 rates as well as on the biogas production. O' Connor et al. <sup>[8]</sup> showed that DEHP  
85 exhibited a relatively high toxicity to methanogenesis over 100 mg/L. It was found in  
86 our lab-scale anaerobic bioreactor effluent at a concentration of 1 mg/L, and its

87 concentration decreased in the aerobic polishing step placed after the anaerobic  
88 bioreactor. <sup>[2]</sup> Jonsson et al. <sup>[9]</sup> found phthalic acids at concentrations of 50 mg/L in  
89 landfills and observed that the concentration decreased over time. Another example of  
90 a commonly found plasticiser is 2,6-di-tert-butyl-4-(dimethylaminomethyl) phenol  
91 (MAMP) which is an antioxidant and stabiliser used as an oxidation inhibitor in  
92 natural and synthetic elastomers, polyolefin plastics, resins, adhesives, petroleum oil  
93 and waxes. <sup>[10]</sup>

94

95 Little is known regarding their biodegradability in the environment and even less  
96 about their toxicity towards methanogens. Methanogens are known to be the most  
97 sensitive trophic group in the anaerobic process, and any inhibition of their  
98 metabolism could cause an anaerobic digester to fail. Anaerobic digestion is  
99 becoming widely used to treat waste, and it is therefore important to know at which  
100 concentration these plasticisers will become an issue. Because some plasticisers can  
101 accumulate in the environment, it is important to know if these could have an effect  
102 on methanogens during the anaerobic treatment of leachate. This is particularly  
103 relevant in the case where plastics or industrial wastes are present in the municipal  
104 solid waste landfill.

105

106

## 107 **MATERIALS AND METHODS**

108

109

110 Three plasticisers (Table 1) that were found to be recalcitrants by GC-MS in our lab-  
111 scale anaerobic process were tested to determine their biodegradability and their

112 toxicity towards methanogens: <sup>[2]</sup> o-hydroxybiphenyl (HBP), Bis(2-  
113 Ethylhexyl)phthalate (DEHP) and 3,5-Di-tert-butyl-4-hydroxyphenyl propionic acid.  
114 Unfortunately, the latter was not commercially available and 2,6-di-tert-butyl-4-  
115 (dimethylaminomethyl) phenol (MAMP) was used instead because of a very similar  
116 structural formula. These plasticisers were purchased from Sigma-Aldrich (analytical  
117 grade).

118 Plasticiser concentrations of 200, 20, 2, 0.2, 0.02 and 0.002 mg/L were tested in  
119 duplicate for each plasticiser in order to determine which concentration would cause  
120 cessation of methanogenesis. A wide range of concentrations was tested in order to  
121 determine not only at which concentration they start to inhibit methanogenesis, but  
122 also determine if at high concentration such as 200 mg/L these can be metabolised  
123 and ultimately converted to methane. Forty mg of acetic acid was put in each 38 mL  
124 glass bottle to act as a carbon source for the methanogens. A total volume of 20 mL of  
125 inoculum, Owen et al.'s buffered biomedium, <sup>[11]</sup> and plasticiser was added to each  
126 bottle while flushing with CO<sub>2</sub>/N<sub>2</sub> gas (30/70) and sealed off immediately after. <sup>[11]</sup>  
127 The inoculum (2 mL in each bottle) was taken from an active anaerobic digester  
128 treating landfill leachate; <sup>[12]</sup> its total suspended solids and volatile suspended solids  
129 content were 22.9 and 16.9 g/L, respectively. Two controls containing acetic acid,  
130 inoculum and biomedium were run in parallel. The glass bottles were incubated in an  
131 orbital shaker at 35°C. The biogas volumes were regularly measured using a wetted  
132 glass syringe and reported at atmospheric pressure and a temperature of 35°C. The  
133 composition of gas was determined using a Shimadzu GC-TCD fitted with a Porapak  
134 N column (1500×6.35 mm).

135

136

137 **RESULTS AND DISCUSSION**

138

139

140 Previous studies on landfill leachate have extensively identified the compounds  
141 present in leachate, but not much is known about the tolerance of methanogens  
142 towards these toxicants. Interference with the metabolism of methanogenic cultures  
143 can manifest itself in several different ways in these tests. If the test compound is  
144 extremely toxic, it may inactivate all the microorganisms responsible for at least one  
145 step in the metabolic sequence. In a slightly less severe situation, the test compound  
146 may totally or partially inhibit microbial metabolism. If the compound does not  
147 completely inhibit metabolism, some bacterial activity will continue, and the culture  
148 may eventually acclimate to the compound, allowing a return to the same specific  
149 metabolic rate as in the absence of toxicant. [7]

150 The ATA was also used to determine whether methane production was due to acetic  
151 acid only, or if the plasticiser was also biodegraded. The evidence of toxicity in an  
152 ATA may be either a decreased initial gas production or a lag phase before gas  
153 production begins. In either case, the toxic effects should diminish with time and the  
154 ultimate gas production will reflect the additional gas generated by utilization of the  
155 test compound. In the bottles with 200 mg/L of plasticiser the theoretical COD masses  
156 introduced were 11.2, 10.3 and 10.5 mg COD for MAMP, DEHP and HBP,  
157 respectively. Hence, if the compounds were biodegraded there should be circa 25 %  
158 additional methane production for the 200 mg/L sample (COD from the plasticiser in  
159 addition to the 40 mg of acetic acid). The methane produced in the control with 40 mg  
160 acetic acid was circa 13 mL which is approximately 80 % of the theoretical value of  
161 0.395 mL at 35°C per g COD removed.

162

163 Figure 1 shows that MAMP was not toxic at concentrations below 20 mg/L. However,  
164 with 200 mg/L, there was partial inhibition as methane production was more sluggish  
165 until day 14 (about half the control on day 14) before it acclimated and then returned  
166 to the same specific rate as in the absence of the plasticiser on day 21. Thus our  
167 results showed that at concentration usually found in leachate (ppb to ppm levels) this  
168 plasticiser will not affect methanogenesis. No additional methane could be produced  
169 from MAMP.

170

171 No inhibitory effects were observed with DEHP (Fig. 2), but DEHP could not be  
172 metabolized to produce methane. DEHP has a high hydrophobicity ( $\log K = 8.7$ )  
173 which make it "stick" to the biomass, however, this did not affect methanogenesis  
174 even at 200 mg/L which is in contradiction with Gavala et al. <sup>[5]</sup> and O'Connor et al.  
175 <sup>[8]</sup> who found severe toxicity at concentrations of 60 and 100 mg/L, respectively. This  
176 shows that landfill leachate containing significant amounts of this persistent  
177 plasticiser could still be treated with no sign of imbalance. Jonsson et al. <sup>[9]</sup> observed  
178 that the concentration of DEHP decreased over time during methanogenesis in a  
179 landfill. This indicated that DEHP can be degraded to phthalic acid directly or via its  
180 monoester, but our results showed that it cannot be ultimately converted to methane.

181

182 In contrast, HBP was found to be the most toxic of the three plasticisers tested (Fig.  
183 3): a severe inhibition was observed at 200 mg/L and it took more than thirty days for  
184 the methanogens to acclimatise and metabolise acetic acid to produce methane gas.  
185 No additional methane was obtained with MAMP and DEHP because of the control  
186 and the 200 mg/L curves were very similar. In contrast, with 200 mg/L of HBP the  
187 methane production was 25 % greater than in the control. Thus, HBP was

188 biodegraded and converted to methane, and the lag phase was an adaptation period  
189 necessary to co-metabolize acetic acid and HBP. This is in agreement with previous  
190 studies where HBP was found to be biodegraded by *Pseudomonas* species via  
191 successive steps. <sup>[13-14]</sup> Thus our results showed that at concentration usually found in  
192 leachate (ppb to ppm levels) these three plasticisers will not affect methanogenesis.

193

194

## 195 **CONCLUSIONS**

196

197 ATA results showed no toxicity effect at concentrations equal to and below 20 mg/L  
198 for MAMP. However, at 200 mg/L a 50 % drop in methane production was observed  
199 on day 14 after which it returned to the same levels as the control. No effects on  
200 methanogens were noticed for Bis(2-Ethylhexyl)phthalate. These two plasticisers may  
201 have been biodegraded but could not be converted to methane. Finally, for o-  
202 hydroxybiphenyl a 75 % drop in methane production was noticed at 20 mg/L on day  
203 14, whereas no gas was produced at 200 mg/L before thirty days. After acclimation,  
204 o-hydroxybiphenyl at 200 mg/L could be biodegraded resulting in a 25 % increase in  
205 methane production compared to the control.

206

## 207 **ACKNOWLEDGMENT**

208

The authors are grateful to the Department of Environment, Food and Rural Affairs  
(DEFRA) in the UK for their sponsorship of this research.

209

## 210 **REFERENCES**

211



- 212 [1] EPA. *Chemicals known to cause cancer or reproductive toxicity*. 2007  
213 [accessed October 2011]; Available from:  
214 [http://www.oehha.ca.gov/prop65/prop65\\_list/files/P65single092807.pdf](http://www.oehha.ca.gov/prop65/prop65_list/files/P65single092807.pdf).
- 215 [2] Trzcinski, A.P.; D.C. Stuckey. Treatment of municipal solid waste leachate  
216 using a submerged anaerobic membrane bioreactor at mesophilic and  
217 psychrophilic temperatures: analysis of recalcitrants in the permeate using  
218 GC-MS. *Water Res.* **2010**, *44*(3), 671-680.
- 219 [3] Yasuhara, A.; H. Shiraishi; M. Nishikawa; T. Yamamoto; O. Nakasugi.  
220 Organic components in leachates from hazardous waste disposal sites. *Waste*  
221 *manag. and res.* **1999**, *17*(3), 186.
- 222 [4] Benotti, M.J.; R.A. Trenholm; B.J. Vanderford; J.C. Holady; B.D. Stanford;  
223 S.A. Snyder. Pharmaceuticals and endocrine disrupting Compounds in U.S.  
224 drinking water. *Environ. Sci. Technol.* **2009**, *43*(3), 597.
- 225 [5] Gavala, H.N.; F. Alatrisme-Mondragon; R. Iranpour; B.K. Ahring.  
226 Biodegradation of phthalate esters during the mesophilic anaerobic digestion  
227 of sludge. *Chemosphere* **2003**, *52*(4), 673-682.
- 228 [6] Bauer, M.J.; R. Herrmann. Estimation of the environmental contamination by  
229 phthalic acid esters leaching from household wastes. *Science of The Total*  
230 *Environment* **1997**, *208*(1-2), 49-57.
- 231 [7] Asakura, H.; T. Matsuto; N. Tanaka. Behavior of endocrine-disrupting  
232 chemicals in leachate from MSW landfill sites in Japan. *Waste Manag.* **2004**,  
233 *24*(6), 613-622.
- 234 [8] O'Connor, O.A.; M.D. Rivera; L.Y. Young. Toxicity and biodegradation of  
235 phthalic acid esters under methanogenic conditions. *Environmental*  
236 *Toxicology and Chemistry* **1989**, *8*, 569-576.

- 237 [9] Jonsson, S.; J. Ejlertsson; B.H. Svensson. Transformation of phthalates in  
238 young landfill cells. *Waste Manag.* **2003**, *23*(7), 641-651.
- 239 [10] Miller, D.; B.B. Wheals; N. Beresford; J.P. Sumpter. Estrogenic activity of  
240 phenolic additives determined by an in vitro yeast bioassay. *Environmental*  
241 *health perspectives* **2001**, *109*(2), 133.
- 242 [11] Owen, W.F.; D.C. Stuckey; J. Healy, J. B.; L.Y. Young; P.L. McCarty.  
243 Bioassay for monitoring biochemical methane potential and anaerobic  
244 toxicity. *Water Res.* **1979**, *13*(6), 485-492.
- 245 [12] Trzcinski, A.P.; D.C. Stuckey. Continuous treatment of the organic fraction of  
246 municipal solid waste in an anaerobic two-stage membrane process with liquid  
247 recycle. *Water Res.* **2009**, *43*(9), 2449-2462.
- 248 [13] Furukawa, K.; N. Arimura; T. Miyazaki. Nucleotide sequence of the 2,3-  
249 dihydroxybiphenyl dioxygenase gene of *Pseudomonas pseudoalcaligenes*. *J.*  
250 *Bacteriol.* **1987**, *169*(1), 427-429.
- 251 [14] Suske, W.A.; W.J.H. van Berkel; H.-P.E. Kohler. Catalytic mechanism of 2-  
252 hydroxybiphenyl 3-monooxygenase, a flavoprotein from *Pseudomonas*  
253 *azelaica* HBP1. *J. Biol. Chem.* **1999**, *274*(47), 33355-33365.
- 254
- 255

256 **LIST OF FIGURE AND TABLE CAPTION**

257

258

259 **Figure 1.** Anaerobic Toxicity Assay of 2,6-di-tert-butyl-4-(dimethylaminomethyl)  
260 phenol (MAMP). The maximum standard deviation was 0.2 mL.

261

262 **Figure 2.** Anaerobic Toxicity Assay of Bis(2-Ethylhexyl)phthalate (DEHP). The  
263 maximum standard deviation was 0.2 mL.

264

265 **Figure 3.** Anaerobic Toxicity Assay of o-hydroxybiphenyl (HBP). The maximum  
266 standard deviation was 0.2 mL.

267

268

269

270

271

272

273

274

275

276

277

278

279


280

281

282

283


284  
285  
286  
287  
288  
289  
290  
291  
292



293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316

Fig. 1


317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334



335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349

Fig. 2

350  
351  
352  
353  
354  
355  
356  
357  
358






359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382

Fig. 3

383  
384  
385  
386  
387  
388

**Table 1.** Properties of the plasticisers used for the ATA test in this study. P = octanol/water partition coefficient. LD<sub>50</sub> = median lethal dose for rats.

Plasticiser (acronym)	o-hydroxybiphenyl (HBP)	Bis(2-Ethylhexyl)phthalate (DEHP)	2,6-di-tert-butyl-4-(dimethylaminomethyl)phenol (MAMP)
Molecular formula	C <sub>12</sub> H <sub>10</sub> O	C <sub>24</sub> H <sub>38</sub> O <sub>4</sub>	C <sub>17</sub> H <sub>29</sub> NO
Structural formula			
CAS number	90-43-7	117-81-7	88-27-7
Molecular weight	170.21	390.56	263.42
LD50 (mg/kg)	1050	1370	343
Density	1.293	0.98	0.95
Log P	2.94	8.7	4.6
Solubility (g/L)	0.26	10 <sup>-4</sup>	6.1 at pH 7

389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399