

bomrang: Fetch Australian Government Bureau of Meteorology Data in R

Adam H Sparks¹, Mark Padgham², Hugh Parsonage³, and Keith Pembleton⁴

¹University of Southern Queensland, Centre for Crop Health, Toowoomba Queensland 4350,
Australia

²University of Salzburg, Inter-Faculty Department of Geoinformatics, 5020 Salzburg, Austria

³Grattan Institute, Carlton Victoria 3053, Australia

⁴University of Southern Queensland, School of Agricultural, Computational and
Environmental Sciences, Toowoomba Queensland 4350, Australia

25 Aug 2017

Paper DOI: <http://dx.doi.org/10.21105/joss.00411>

Software Repository: <https://github.com/ropensci/bomrang>

Software Archive: <http://dx.doi.org/10.5281/zenodo.897183>

Summary

The Australian Bureau of Meteorology (BoM) publicly provides data via an anonymous FTP server, in XML and json files (Australian Bureau of Meteorology 2017). The files are well structured but knowledge of how to use R and parse them is required to extract the data into a data frame for use in R (R Core Team 2017) or requires external programs and scripting to import the data for use. *bomrang* provides functionality for automated retrieval and parsing of selected weather data files from BoM. Data that can be fetched include current weather at a given station, daily précis (short summaries less than 30 characters) forecasts for all Australian forecast locations, agricultural bulletins summarising weather observations useful for agriculture for each state or territory and satellite imagery in GeoTIFF file formats. A family of functions, `get_current_weather()`, `get_precis_forecast()`, `get_ag_bulletin()` and `get_weather_bulletin()`, provide the ability to easily download data from BoM, import it and create a tidy data frame (Wickham 2014) of the data. To help identify stations given a specific location, `sweep_for_station()` returns a data frame of all weather stations (in this package) sorted by distance from a user specified latitude and longitude, ascending. Two further functions, `update_forecast_towns()` and `update_station_locations()`, provide the user with the ability to update internal databases of forecast locations, station metadata and JSON URLs used by the package to interface with BoM. The package's internal databases decrease the time necessary to gather forecast information and return a data frame, while rarely changing. Functionality for automated downloading and importing of satellite imagery is provided by the `get_satellite_imagery()` function. A helper function, `get_available_imagery()`, returns values of currently available satellite imagery for download.

The data have many applications. In agriculture the data are used in several types of models, some of which include the estimation of surface moisture, crop yield estimates, crop development stages or stress or forecasting of epidemics of crop diseases or populations of insect pests [Venäläinen and Heikinheimo (2002); De Wolf, Madden, and Lipps (2003); Sparks2017]. Other areas of use include mapping potential renewable energy, *e.g.* wind or solar potential for exploration purposes (Ramachandra and Shruthi 2007). The data can

also be used by decision makers for municipalities to help planning for extreme weather events, energy needs and other infrastructure (Svensson and Eliasson 2002; Alcoforado et al. 2009).

References

- Alcoforado, Maria-João, Henrique Andrade, António Lopes, and João Vasconcelos. 2009. “Application of Climatic Guidelines to Urban Planning.” *Landscape and Urban Planning* 90 (1): 56–65. doi:<http://dx.doi.org/10.1016/j.landurbplan.2008.10.006>.
- Australian Bureau of Meteorology. 2017. “Weather Data Services.” <http://www.bom.gov.au/catalogue/data-feeds.shtml>.
- De Wolf, E. D., L. V. Madden, and P. E. Lipps. 2003. “Risk Assessment Models for Wheat Fusarium Head Blight Epidemics Based on Within-Season Weather Data.” *Phytopathology* 93 (4). Scientific Societies: 428–35. doi:[10.1094/PHYTO.2003.93.4.428](https://doi.org/10.1094/PHYTO.2003.93.4.428).
- R Core Team. 2017. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Ramachandra, T.V., and B.V. Shruthi. 2007. “Spatial Mapping of Renewable Energy Potential.” *Renewable and Sustainable Energy Reviews* 11 (7): 1460–80. doi:<http://dx.doi.org/10.1016/j.rser.2005.12.002>.
- Svensson, Marie K., and Ingegärd Eliasson. 2002. “Diurnal Air Temperatures in Built-up Areas in Relation to Urban Planning.” *Landscape and Urban Planning* 61 (1): 37–54. doi:[http://dx.doi.org/10.1016/S0169-2046\(02\)00076-2](http://dx.doi.org/10.1016/S0169-2046(02)00076-2).
- Venäläinen, Ari, and Martti Heikinheimo. 2002. “Meteorological Data for Agricultural Applications.” *Physics and Chemistry of the Earth, Parts A/B/C* 27 (23): 1045–50. doi:[http://dx.doi.org/10.1016/S1474-7065\(02\)00140-7](http://dx.doi.org/10.1016/S1474-7065(02)00140-7).
- Wickham, Hadley. 2014. “Tidy Data.” *Journal of Statistical Software* 59 (1): 1–23. doi:[10.18637/jss.v059.i10](https://doi.org/10.18637/jss.v059.i10).