

## Association between Physical Activity and Diabetic Complications among Bangladeshi Type 2 Diabetic Patients

*MS Bukht<sup>1</sup>; KR Ahmed<sup>2,3</sup>; S Hossain<sup>4</sup>; P Masud<sup>5</sup>; S Sultana<sup>2</sup>; R Khanam<sup>6</sup>*

*<sup>1</sup>Health, Nutrition and Population Programme, BRAC, Dhaka, Bangladesh; <sup>2</sup>Dept of Health Promotion and Health Education, Bangladesh University of Health Sciences; <sup>3</sup>School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Australia; <sup>4</sup>Dept of Public Health and Informatics, Jahangirnagar University, Savar, Dhaka, Bangladesh; <sup>5</sup>Aureus University School of Medicine, Aruba, USA; <sup>6</sup>School of Commerce, University of Southern Queensland, Toowoomba, Australia*

### **Abstract:**

**Background:** Physical activity (PA) is a cornerstone of type 2 diabetes management, but is often underutilized. The aim of the study was to assess the association of PA with the presence of diabetes related complications among Bangladeshi type 2 diabetic subjects.

**Methods:** This cross-sectional study was conducted in the OPD of BIRDEM, a tertiary diabetic care hospital in Bangladesh. A group of 977 subjects were randomly selected and followed up. A structured questionnaire was administered to collect socio-demographic information. Diabetes was diagnosed following the WHO study group criteria. The level of PA was categorized into inactive/low (<150 min/week) and moderate-to-vigorous ( $\geq 150$  min/week). The WHO recommended Asian criteria was used to identify general obesity as healthy weight (BMI 18.5–22.99 kg/m<sup>2</sup>), overweight (23.0–24.99 kg/m<sup>2</sup>) and obese (25.0 and above kg/m<sup>2</sup>). Retinopathy was detected by fundal photography, CKD by serum creatinine and hypertension was diagnosed clinically. Univariate and multivariate generalized linear models were used to assess the associations of physical activity with diabetes related complications.

**Results:** Out of the 977 subjects investigated, 468 were male and 509 were female (mean±SD of age, 56±8 years). In the study subjects, 74% were either inactive or in low PA and of them 65.1% were overweight/obese. Analysis showed that inactive/low PA was associated with all the three diabetes related complications ( $p<0.001$ )—hypertension, retinopathy, and nephropathy. Multivariate analysis showed that inactive/low level of PA was strongly associated with complications like retinopathy ( $p<0.001$ ) and hypertension ( $p=0.01$ ) in the female patients. It was also found to be highly associated with retinopathy ( $p <0.001$ ) among the male patients.

**Conclusion:** A large number of urban of Bangladeshi population are involved only in low PA or remain inactive themselves, which is leading to obesity and it seems to have a strong association with diabetes related complications in this population.

**Key words:** Physical activity, Type 2 diabetes, Hypertension, Diabetic retinopathy, Diabetic nephropathy.

**Introduction:**

Diabetes mellitus (DM) has emerged as a major public health challenge around the world. Not only the developed country but also low and middle-income countries face the greatest burden of DM. It has become one of the major causes of mortality in the long run and based on WHO fact sheet, an estimated 1.6 million deaths were directly caused by diabetes globally in year 2015 (WHO 2017). The global prevalence of diabetes among adults over 18 years of age has increased and was recorded to be 8.5% in year 2014 (WHO 2017). In developed countries, diabetes is known to be the seventh leading cause of death in year 2015 (CDC 2017). The Diabetes Atlas of the International Diabetes Federation (IDF) estimated in 2011 that the global DM prevalence was 8.3% in the age group 20–79 years. The number of people living with DM is projected to reach 551.9 million by 2030 (Whiting et al.). Bangladesh is one of the countries out of many in South Asian region which has the second largest number of adults with diabetes; more than 7.1 million, 8.4% or 10 million according to research published in WHO bulletin in 2013 (Mahbub 2016). By 2030, Bangladesh is likely to emerge as the eighth highest-ranking country in terms of the number of people with DM (Whiting et al. 2011).

The management of diabetes mellitus and the management and prevention of the complications are important challenges of the present time. Studies indicate that genetic factors do not account entirely for the development of diabetes, and several environmental triggers have been implicated (Akerblom et al. 2002). Physical activity is the key element in the prevention and management of type 2 diabetes. For patients with diabetes, insufficient physical activity is considered the most important risk factors for developing macro and microvascular complications (ADA 2003). Regular physical

activity helps to maintain weight loss and prevent weight regain (Wing and Hill 2001). Regular exercise and aerobic fitness also improve insulin sensitivity, glycaemic control (Bird and Hawley 2016; Way et al. 2016), may decrease the risk of developing diabetes (Canadian Diabetes Association Clinical Practice Guidelines Expert et al. 2013; Wei M et al. 1999), and may reduce overall mortality in patients who have type 2 diabetes (Kokkinos et al. 2009). Physical activity exerts numerous beneficial health effects on glycaemic control, insulin sensitivity, blood pressure, lipid profile, and endothelial function, and the evidence favouring a physically active lifestyle in the treatment of chronic diseases is substantial (Pedersen and Saltin 2015). With the help of many studies that has been conducted over the years, it is now well established that participation in regular physical activity can improve blood glucose control and can prevent or delay type 2 diabetes, along with positively affecting lipids, blood pressure, cardiovascular events, mortality, and quality of life (Wahid et al. 2016).

However, studies have shown that about 35% of diabetic patients from Bangladesh do not participate in any physical activity (Saleh et al. 2012). For Bangladeshi diabetic patients, this could be followed for the elevation of the complications of the disease. Before that, the fact we need to evaluate the scenario of the association between physical activity and the diabetic complications are extremely needed for Bangladeshi diabetic patients. Moreover, there is still a paucity of published studies on this association among Bangladeshi type 2 diabetic mellitus (T2DM) patients. Therefore, the aim of this study was to investigate the association of physical activity with the presence of diabetes related complications among Bangladeshi type 2 diabetic patients.

## ***Methods***

### ***Study design, and setting***

A hospital-based cross sectional study was conducted in the years of 2009 and 2010 among a representative sample of Bangladeshi T2DM patients. A total, 977 patients were randomly selected from the Out-Patient Department (OPD) of the Bangladesh Institute of Research and Rehabilitation in Diabetes, Endocrine and Metabolic Disorders (BIRDEM), which serves as the primary destination for diabetic patient across Bangladesh. Furthermore, BIRDEM is a sister concern of the Diabetic Association of Bangladesh and is a WHO Collaborating Centre for research into the prevention and control of diabetes. In addition to having its own primary care facility, BIRDEM serves as a referral center for 57 affiliated associations. Being a charity organization, it attracts patients from diverse demographic and socio-economic backgrounds from across Bangladesh. Routinely, newly diagnosed patients in the BIRDEM undergo thorough clinical and biochemical investigations to determine their diabetic profile and the presence of any complications.

### ***Study procedures***

Data were collected in a predesigned interviewer administered questionnaire. The selection of the study participants was based on some inclusion and exclusion criteria. Patients, aged  $\geq 30$ , with type 2 diabetes having necessary clinical reports were primarily included in the sampling list. This list was further refreshed by excluding the patients

with pregnancy, advised for hospitalization, and those who were unable to answer to a short list of simple questions about their disease and PA.

**Measures and statistical analysis:**

Physical and anthropometric measurements were taken from the patients adopting standard operating procedure. The classification and diagnosis of diabetes was made by the reporting physician according to the World Health Organization (WHO) criteria (Alberti and Zimmet 1998). The ophthalmologic examination included best-corrected visual acuity with an illuminated Snellen chart, slit-lamp bio microscopy and also detailed fundus examination by both indirect ophthalmoscopy and contact lens bio-microscopy conducted by retinal specialist (Ahmed et al. 2011). Retinopathy was categorized from fundus photography, CKD by serum creatinine, and we used the grading categories of retinopathy as no diabetic retinopathy (NDR) and any stages of diabetic retinopathy (DR). The cutoff considered for hypertension was blood pressure  $\geq 140/\geq 90$  mmHg. Patients on antihypertensive drugs have also been considered as hypertensive. In this study, the level of PA was categorized into inactive/low ( $<150$  min/week) and moderate-to-vigorous ( $\geq 150$  min/week) in accordance with the American College of Sports Medicine and the American Diabetes Association (Colberg et al. 2010). The WHO recommended Asian criteria was used to identify general obesity as healthy weight (BMI 18.5–22.99 kg/m<sup>2</sup>), overweight (23.0–24.99kg/m<sup>2</sup>) and obese (25.0 and above kg/m<sup>2</sup>) (Hossain et al. 2018; WHO 2004).

**Statistical analysis:**

All statistical analyses were performed by using Statistical Package for Social Science (SPSS) version 20. The descriptive statistics of mean, standard deviation and percentages were calculated for the entire sample. Chi-square test was used to examine differences in the prevalence of different categorical variables. Both univariate and multivariate analysis were performed to assess the associations of physical activity with diabetes related complications. The 'p' value less than 0.05 was considered as statistically significant for this study.

**Ethical issues:**

This study was conducted maintaining ethical standards to the highest possible extent. Prior to the assessment, informed consent was taken from all the patients, participated in this study. This study was approved by the Ethics Committees of the Diabetic Association of Bangladesh.

**Results:**

A total of 977 diabetic patients were interviewed (Table 1) of which 468 were male (47.9%) and 509 were female (52.1%). A plurality of the study participants' age range was between 46 and 60 years (63.2%) and the mean age of all the participants was  $56 \pm 8$  years. Majority of the patients were obese (35.5%). The percentage of both overweight and obesity was 64.1%. Approximately, one fourth of the participants (26%) were involved in moderate-to-vigorous PA, where females were more active (68.1%) than their

male counterpart, which is statistically highly significant ( $p < 0.001$ ; Table 2). Out of 723 diabetic patients who had inactive/low PA, 65.1% were overweight/obese.

Irrespective of gender, this study revealed that inactive/low PA was significantly associated with all the three diabetes related complications—hypertension ( $p < 0.001$ ), retinopathy ( $p < 0.001$ ), and nephropathy ( $p < 0.001$ ) (Table 3). Binary logistic regression analysis showed that the patients who were inactive or involved in low level of PA, males were 21 times (95% CI: 8.55–52.55) and female patients were 22 times (95% CI: 12.03–40.51) higher likely to suffer from diabetic retinopathy (Table 4). In terms of female diabetic patients, it was also found that inactive/low PA was strongly associated with the complication, hypertension ( $p = 0.010$ ).

### **Description:**

To the best of our knowledge, this is the first study to report Bangladeshi population-level estimates of the association between a range of diabetes complications and PA. In agreement with the previous epidemiological studies (Hjerkind et al. 2017; Jerums and MacIsaac 2015; Lee et al. 2018; Medina et al. 2018; Nguyen et al. 2017; Praidou et al. 2017), we found that inactive/low level of PA was statistically highly significant with hypertension, retinopathy and nephropathy of the type 2 diabetic patients. Inactive in PA and/or low levels of PA is independent, modifiable risk factors for the progression of insulin resistance, and poor health outcomes with type 2 diabetes (Booth et al. 2012; Van Buren and Tibbs 2014). Our study showed that female diabetic patients were more vulnerable to the type 2 diabetic complications such as hypertension and retinopathy than

their male counterpart was. Similar findings were also found among the type 2 diabetic patients in middle-aged population in Pakistan (Ansari 2009).

Physical inactivity, a preventable risk factor for obesity that is significantly associated with diabetes mellitus (Hjerkind et al. 2017). The upsurge in the prevalence of type 2 diabetes is inextricably linked to the increase in obesity (Verma and Hussain 2017). Our study found that nearly two thirds of the diabetic patients with low PA in Bangladesh were obese. This finding justify the findings a review report of Eckel et. al. (Eckel et al. 2011). Recent systematic reviews on intervention studies, performed in different settings and populations, proved that physical activity focused lifestyle interventions can be effective in prevention of hypertension (Liu et al. 2017), diabetes mellitus and its complications (Asano et al. 2014; Balducci et al. 2009; Janevic et al. 2013). It is proved by the systematic review and meta-analysis that observance to guidelines suggested minimum physical activity level reduced incidence of both hypertension (RR = 0.94; 95% CI: 0.92–0.97) and diabetes mellitus (RR = 0.74, 95% CI: 0.72–0.77) (Asano et al. 2014; Liu et al. 2017; Wahid et al. 2016).

### **Conclusions:**

The principle implications of this study is to highlight the importance of physical activity for the prevention of type 2 diabetes in the middle-aged population of Bangladesh which is experiencing a rapid and substantial decline of physical activity levels as a result of poor dietary habits and sedentary behaviors. In parallel with decreasing levels of physical activity, the prevalence of obesity has increased significantly in Bangladesh and

consequently, diabetes mellitus has become a major public health issue. Therefore, promoting an active lifestyle or regular exercise should receive the highest public health priority in the country. A large number of urban of Bangladeshi population are involved in only low level of PA, which is leading to obesity, and it seems to have a strong association with diabetes related complications in this population. Furthermore, coordinated education and motivation program should be undertaken to improve awareness on this issue.

#### **References:**

- ADA (2003) Physical Activity/Exercise and Diabetes Mellitus. American Diabetes Association. *Diabetes care* 26(Supplement 1):S73-S77  
doi:10.2337/diacare.26.2007.S73
- Ahmed KR, et al. (2011) Risk factors of diabetic retinopathy in Bangladeshi type 2 diabetic patients. *Diabetes & metabolic syndrome* 5(4):196-200  
doi:10.1016/j.dsx.2012.02.020
- Akerblom HK, Vaarala O, Hyoty H, Ilonen J, Knip M (2002) Environmental factors in the etiology of type 1 diabetes. *American journal of medical genetics* 115(1):18-29  
doi:10.1002/ajmg.10340
- Alberti KGMM, Zimmet PZ (1998) Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus. Provisional report of a WHO Consultation. *Diabetic Medicine* 15(7):539-553  
doi:10.1002/(sici)1096-9136(199807)15:7<539::aid-dia668>3.0.co;2-s
- Ansari RM (2009) Effect of physical activity and obesity on type 2 diabetes in a middle-aged population. *Journal of environmental and public health* 2009:195285  
doi:10.1155/2009/195285
- Asano RY, et al. (2014) Acute effects of physical exercise in type 2 diabetes: A review. *World journal of diabetes* 5(5):659-65  
doi:10.4239/wjd.v5.i5.659
- Balducci S, et al. (2009) Physical activity/exercise training in type 2 diabetes. The role of the Italian Diabetes and Exercise Study. *Diabetes/metabolism research and reviews* 25 Suppl 1:S29-33  
doi:10.1002/dmrr.985
- Bird SR, Hawley JA (2016) Update on the effects of physical activity on insulin sensitivity in humans. *BMJ open sport & exercise medicine* 2(1):e000143  
doi:10.1136/bmjsem-2016-000143

- Booth FW, Roberts CK, Laye MJ (2012) Lack of exercise is a major cause of chronic diseases. *Comprehensive Physiology* 2(2):1143-211 doi:10.1002/cphy.c110025
- Canadian Diabetes Association Clinical Practice Guidelines Expert C, et al. (2013) Physical activity and diabetes. *Canadian journal of diabetes* 37 Suppl 1:S40-4 doi:10.1016/j.cjcd.2013.01.018
- CDC (2017) New CDC report: More than 100 million Americans have diabetes or prediabetes. Centers for Disease Control and Prevention. <https://www.cdc.gov/media/releases/2017/p0718-diabetes-report.html>. (Access date: July 10, 2018)
- Colberg SR, et al. (2010) Exercise and type 2 diabetes: the American College of Sports Medicine and the American Diabetes Association: joint position statement. *Diabetes care* 33(12):e147-67 doi:10.2337/dc10-9990
- Eckel RH, et al. (2011) Obesity and type 2 diabetes: what can be unified and what needs to be individualized? *The Journal of clinical endocrinology and metabolism* 96(6):1654-63 doi:10.1210/jc.2011-0585
- Hjerkind KV, Stenehjem JS, Nilsen TI (2017) Adiposity, physical activity and risk of diabetes mellitus: prospective data from the population-based HUNT study, Norway. *BMJ open* 7(1):e013142 doi:10.1136/bmjopen-2016-013142
- Hossain S, Hossain MS, Anjum A, Ahmed F, Hossain MF, Uddin ME (2018) Risk modeling of non-communicable diseases using socio-demographic characteristics, lifestyle and family disease history among university students in Bangladesh. *Journal of Public Health* doi:10.1007/s10389-018-0895-7
- Janevic MR, McLaughlin SJ, Connell CM (2013) The association of diabetes complications with physical activity in a representative sample of older adults in the United States. *Chronic illness* 9(4):251-7 doi:10.1177/1742395313475461
- Jerums G, MacIsaac RJ (2015) Diabetic nephropathy: How does exercise affect kidney disease in T1DM? *Nature reviews Endocrinology* 11(6):324-5 doi:10.1038/nrendo.2015.46
- Kokkinos P, et al. (2009) Exercise capacity and all-cause mortality in African American and Caucasian men with type 2 diabetes. *Diabetes care* 32(4):623-8 doi:10.2337/dc08-1876
- Lee JY, Ryu S, Sung KC (2018) Association of baseline level of physical activity and its temporal changes with incident hypertension and diabetes mellitus. *European journal of preventive cardiology* 25(10):1065-1073 doi:10.1177/2047487318774419
- Liu X, et al. (2017) Dose-Response Association Between Physical Activity and Incident Hypertension: A Systematic Review and Meta-Analysis of Cohort Studies. *Hypertension* 69(5):813-820 doi:10.1161/HYPERTENSIONAHA.116.08994
- Mahbub I (2016) The State Of Diabetes In Bangladesh. *Beta Future Startup*. <https://futurestartup.com/2016/07/27/the-state-of-diabetes-in-bangladesh/>. (Access Date: July 10, 2018)
- Medina C, Janssen I, Barquera S, Bautista-Arredondo S, Gonzalez ME, Gonzalez C (2018) Occupational and leisure time physical inactivity and the risk of type II diabetes and hypertension among Mexican adults: A prospective cohort study. *Scientific reports* 8(1):5399 doi:10.1038/s41598-018-23553-6

- Nguyen B, Bauman A, Ding D (2017) Incident Type 2 Diabetes in a Large Australian Cohort Study: Does Physical Activity or Sitting Time Alter the Risk Associated With Body Mass Index? *Journal of physical activity & health* 14(1):13-19 doi:10.1123/jpah.2016-0184
- Pedersen BK, Saltin B (2015) Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scandinavian journal of medicine & science in sports* 25 Suppl 3:1-72 doi:10.1111/sms.12581
- Praidou A, Harris M, Niakas D, Labiris G (2017) Physical activity and its correlation to diabetic retinopathy. *Journal of diabetes and its complications* 31(2):456-461 doi:10.1016/j.jdiacomp.2016.06.027
- Saleh F, Mumu SJ, Ara F, Ali L, Hossain S, Ahmed KR (2012) Knowledge, Attitude and Practice of Type 2 Diabetic Patients Regarding Obesity: Study in a Tertiary Care Hospital in Bangladesh. *Journal of public health in Africa* 3(1):e8 doi:10.4081/jphia.2012.e8
- Van Buren DJ, Tibbs TL (2014) Lifestyle interventions to reduce diabetes and cardiovascular disease risk among children. *Current diabetes reports* 14(12):557 doi:10.1007/s11892-014-0557-2
- Verma S, Hussain ME (2017) Obesity and diabetes: An update. *Diabetes & metabolic syndrome* 11(1):73-79 doi:10.1016/j.dsx.2016.06.017
- Wahid A, et al. (2016) Quantifying the Association Between Physical Activity and Cardiovascular Disease and Diabetes: A Systematic Review and Meta-Analysis. *Journal of the American Heart Association* 5(9) doi:10.1161/JAHA.115.002495
- Way KL, Hackett DA, Baker MK, Johnson NA (2016) The Effect of Regular Exercise on Insulin Sensitivity in Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis. *Diabetes & metabolism journal* 40(4):253-71 doi:10.4093/dmj.2016.40.4.253
- Wei M, Gibbons LW, Mitchell TL, Kampert JB, Lee CD, SN B (1999) The association between cardiorespiratory fitness and impaired fasting glucose and type 2 diabetes mellitus in men. *Ann Intern Med* 130(2):89-96
- Whiting DR, Guariguata L, Weil C, Shaw J (2011) IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes research and clinical practice* 94(3):311-21 doi:10.1016/j.diabres.2011.10.029
- WHO (2004) Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet* 363(9403):157-163 doi:10.1016/s0140-6736(03)15268-3
- WHO (2017) Fact Sheets on Diabetes. World Health Organizations. <http://www.who.int/news-room/fact-sheets/detail/diabetes>. (Access date: July 10, 2018)
- Wing RR, Hill JO (2001) Successful weight loss maintenance. *Annual review of nutrition* 21:323-41 doi:10.1146/annurev.nutr.21.1.323

**Table 1:** Sample characteristics (n=977)

<b>Variable</b>	<b>Result, n (%)</b>
Gender	
Male	468 (47.9)
Female	509 (52.1)
Age Group	
<45 years	115 (11.8)
46-60 years	617 (63.2)
>60 years	245 (25.1)
Employment status	
Employed	373 (38.2)
Unemployed	604 (61.8)
Physical Activity Level	
Inactive/low	723 (74)
Moderate to vigorous	254 (26)
BMI category	
Underweight	19 (1.9)
Normal weight	332 (34)
Overweight	279 (28.6)
Obese	347 (35.5)
Mean of the variables	
Age in years (m $\pm$ SD)	56 $\pm$ 8
Monthly family income	20026.10 $\pm$ 12534.08
BMI (m $\pm$ SD)	24.18 $\pm$ 2.95

Results are expressed as mean (m)  $\pm$  standard deviation (SD) and number (n) of the respondents. *BMI* Body Mass Index.

**Table 2:** Distribution of PA level according to age, gender and BMI category (n=977)

Variables	Level of physical activity		$\chi^2$ value (p-value)**
	Inactive/low, n (%)	Moderate to vigorous, n (%)	
Gender			
Male	387 (53.5)	81 (31.9)	35.26 (<0.001)
Female	336 (46.5)	173 (68.1)	
Age Group			
<45 years	69 (9.5)	46 (18.1)	79.09 (<0.001)
46-60 years	421 (58.2)	196 (77.2)	
>60 years	233 (32.3)	12 (4.7)	
Overweight and/or obesity			
Yes	471 (65.1)	155 (61.0)	1.39 (0.24)
No	252 (34.9)	99 (39.0)	

\*\* p value is significant at the 0.01 level (2-tailed)

**Table 3:** Relationship between inactive/low PA level and diabetic complications according to gender (n=977)

<b>Complications</b>	<b>Inactive/low physical activity</b>	
	<b>Male (r/p values)</b>	<b>Female (r/p values)</b>
Hypertension	0.162/<0.001	0.252/<0.001
Retinopathy	0.414/<0.001	0.206/<0.001
Nephropathy	0.175/<0.001	0.569/<0.001

*Spearman's non-parametric correlation coefficient's test was done as a test of significance. Correlation is significant at the 0.01 level (2-tailed)*

**Table 4:** Association of physical activity with various complications of the diabetic patients as explored by binary logistic regression (n=977)

Complications	$\beta$ Value	p value	Exp(B)	95.0% CI	
				Lower Bound	Upper Bound
<i>Male</i>					
Hypertension	0.010	0.973	1.010	0.584	1.746
Retinopathy	3.054	<b>&lt;0.001</b>	21.200	8.553	52.550
Nephropathy	0.101	0.718	0.904	0.522	1.565
<i>Female</i>					
Hypertension	0.610	<b>0.010</b>	1.841	1.159	2.925
Retinopathy	3.095	<b>&lt;0.001</b>	22.079	12.034	40.505
Nephropathy	0.298	0.232	0.742	0.455	1.210

*$\beta$  for standardized regression coefficient. Physical activity was taken as dependent variable, whereas other variables were taken as independent variable.*