

What do pedometer counts represent? A comparison between pedometer data and data from four different questionnaires

Katrien A De Cocker*, Ilse M De Bourdeaudhuij and Greet M Cardon

Department of Movement and Sports Sciences, Ghent University, Watersportlaan 2, B-9000 Ghent, Belgium

Submitted 12 June 2007; Accepted 14 January 2008; First published online 20 March 2008

Abstract

Objectives: To compare physical activity (PA) reported through pedometer registrations (step counts) with PA reported in four different questionnaires; to compare step count thresholds (7500, 10 000 and 12 500 steps/d) with the PA guideline of 30 min of moderate to vigorous PA (MVPA) per day.

Subjects: A sample of 310 healthy adults, mean age 38·7 (SD 11·9) years, volunteered to participate. Forty-seven per cent was male and 93% of the sample was employed.

Methods: PA was assessed by interview (Minnesota Leisure Time Physical Activity Questionnaire (MLTPAQ)), three self-administered questionnaires (long version and short version of the International Physical Activity Questionnaire (long-form IPAQ, short-form IPAQ), Baecke questionnaire) and seven consecutive days of pedometer registration.

Results: Step counts correlated positively with questionnaire-based PA. The strongest correlations were found between step counts and total PA reported in the long-form IPAQ ($r_s = 0\cdot37$), moderate PA reported in the short-form IPAQ ($r_s = 0\cdot33$), total and moderate PA reported in the MLTPAQ ($r_s = 0\cdot32$), and the total and leisure-time PA indices (excluding sport) reported in the Baecke questionnaire ($r_s = 0\cdot44$). According to step counts, 22·6% of the participants were somewhat active, 18·7% active and 39·4% highly active. As assessed by the long-form IPAQ, short-form IPAQ and MLTPAQ, the guideline of 30 min MVPA/d was reached by respectively 85·4%, 84·8% and 68·0% of participants.

Conclusion: Pedometer-based data offer adequate information to discriminate between levels of PA. Caution is needed when comparing active samples based on different PA recommendations.

Keywords
Step counter
Survey
Health guidelines
IPAQ

The amount of usual physical activity (PA) accumulated by an individual is strongly related with all-cause morbidity and mortality risk⁽¹⁾. Healthy adults should accumulate 30 min or more of moderate-intensity aerobic PA on 5 d each week or a minimum of 20 min of vigorous-intensity aerobic activity on 3 d each week⁽²⁾. The use of step count recommendations as useful behavioural PA targets has been suggested in the past^(3,4). These step count thresholds are more practical than the 30 min MVPA/d guideline since they do not imply constant time tracking and summing of at least moderate-intensity activity during the day, which is impractical for PA assessment on individual and population levels. Additionally, more detailed pedometer-based indices for public health (sedentary: <5000 steps/d; low active: 5000–7499 steps/d; somewhat active: 7500–9999 steps/d; active: 10 000–12 499 steps/d; highly active: $\geq 12 500$ steps/d) have been introduced by Tudor-Locke and Bassett⁽⁵⁾.

Being able to accurately quantify the amount of PA accrued during daily life is necessary for researchers and health professionals to better understand PA behaviour and to develop successful programmes to increase activity levels in various populations. Traditionally, PA has been measured through questionnaires, the strengths and limitations of which are well known⁽⁶⁾. Recognized benefits are the possibility to assess different dimensions of PA, the ability to collect data from a large number of people at low cost, and the unobtrusiveness of the instrument⁽⁷⁾. However, there are limitations in subjects' recall ability and social desirability bias can cause over-reporting of PA⁽⁷⁾. Recently, more attention has been given to objective instruments to assess PA, e.g. accelerometers and pedometers. Pedometers have been shown to provide a valid and accurate measure of ambulatory activities in free-living conditions^(8,9). They are simple to use, more objective than surveys, and less expensive (approximately \$US 20–50) compared with

*Corresponding author. Email Katrien.DeCocker@UGent.be

accelerometers (approximately \$US 150–500). Despite their limitations such as variability between different brands, insensitivity to non-ambulatory activities (i.e. cycling, swimming) and increased errors at slow walking speeds (<60 m/min)⁽⁸⁾, pedometers have become popular devices for public use and for researchers. Moreover, step counts have proved to be useful in PA studies in free-living populations⁽⁸⁾. However, the question remains of which exactly dimensions of PA are measured by pedometers. Several researchers^(8,10) report that pedometer step counts are an appropriate measurement of the distance walked, while others believe that pedometers also enable the quantification of ambulatory behaviour during occupational, leisure-time, household and transportation activity^(11,12). In addition, one could wonder if step counts are simply a measure for walking or if they can also be associated with self-reports of PA that encompass intensity (other moderate and vigorous PA, besides walking). Therefore, the first purpose of the present study was to compare pedometer data with data from questionnaires, the most commonly used⁽⁷⁾ and practical method for PA assessment in large-scale studies⁽¹³⁾. The objective was to evaluate the associations between step counts and PA (walking, moderate PA, vigorous PA) reported in four different validated questionnaires: the interviewer-administered Minnesota Leisure Time Physical Activity Questionnaire⁽¹⁴⁾ (MLTPAQ), the self-administered long version and self-administered short version of the International Physical Activity Questionnaire⁽¹⁵⁾ (long-form IPAQ, short-form IPAQ), and the self-administered Baecke Questionnaire⁽¹⁶⁾. Additionally, questionnaire-based PA was compared between the five 'step count groups' based on the pedometer health indices of Tudor-Locke and Bassett⁽⁵⁾.

A third objective of the present study was to compare step count thresholds (7500, 10 000 and 12 500 steps/d) with the guideline of 30 min of moderate to vigorous PA (MVPA) per day. Wilde *et al.*⁽¹⁷⁾ found that sedentary women who added a 30 min walk to their daily habits accumulated approximately 10 000 steps. Another study⁽¹⁸⁾ also revealed that on days when women took a 30 min walk, their average step count was close to 10 000. Other researchers⁽¹⁹⁾ found that women who took 10 000 steps/d were more likely to meet the current MVPA guideline, compared with those not accumulating as many steps.

Miller and Brown⁽²⁰⁾ found only a moderate level of agreement between meeting 10 000 steps/d and 150 min or more of PA over five or more sessions per week. However, no research could be located in either gender or in a European sample assessing whether step count thresholds are corresponding to the 30 min MVPA/d guideline. Therefore, the present study explored whether adults reaching 7500, 10 000 and 12 500 steps/d also reached 30 min MVPA/d, based on self-reports.

Methods

Participants and procedures

A convenience sample of 310 volunteers (146 men) living in Flanders, Belgium participated in the present study. Participant characteristics are shown in Table 1. No gender differences were found for the proportions of employed participants, mean ages and average daily step counts.

All participants were visited at home by research assistants for the completion of the MLTPAQ. After this interview, participants were asked to complete the self-administered long-form IPAQ, the self-administered short-form IPAQ and the Baecke questionnaire. Then, participants were asked to register pedometer-based step counts for seven consecutive days. They were instructed to wear the pedometer on their waistband or belt during waking hours. All participants were asked to carry on their usual activities, to remove the pedometer only while swimming, bathing or showering, and to complete an activity log at the end of each day. All participants provided written informed consent and the study was approved by the Ethical Committee of Ghent University. Research assistants (masters-level graduates, native speakers) were given 2 h of training and interview practice.

Instruments

Pedometers

The Yamax Digiwalker SW-200 (Yamax, Tokyo, Japan) was used in the present study as it is known to be a valid, reliable and accurate instrument for counting steps in adult populations⁽²¹⁾.

Table 1 Characteristics of the study participants: healthy adult volunteers, Flanders, Belgium

	Total group		Male		Female		Gender comparison
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
Gender	310	100.0	146	47.1	164	52.9	$\chi^2 = 1.05^{NS}$
Employed	288	92.9	139	95.2	149	90.9	$\chi^2 = 0.35^{NS}$
	Mean	SD	Mean	SD	Mean	SD	
Age (years)	38.7	11.9	39.1	12.5	38.3	11.4	$t = 0.56^{NS}$
Daily step counts	12 087	4888	12 452	4983	11 762	4792	$t = 1.24^{NS}$

NS, $P > 0.05$.

Activity log

Based on procedures of Tudor-Locke *et al.*⁽²²⁾, participants were asked to keep daily activity records in an activity log during the seven consecutive days of pedometer registration. They were requested to record the date, the number of steps recorded at the end of the day, and the type and duration of non-ambulatory activities (e.g. 20 min of biking/swimming). Following established guidelines^(23,24), 150 steps were added to the daily total for every minute of reported biking and/or swimming. The average daily step count was 9981 (SD 3455) without adding the equivalent for biking/swimming and 12 087 (SD 4888) with the added steps for biking/swimming. Of the total sample, 152 people (49.0%) reported biking/swimming during the week of pedometer registration (mean: 7.6 (SD 15.1) min/d).

MLTPAQ

A structured interview was used to solicit detailed information on leisure-time PA over a 1-year period. A Dutch version of the validated MLTPAQ, developed for the Belgian Physical Fitness Study, was used^(6,14,25). Participants' engagement in different activities was queried, together with the number of months per year, the monthly frequency and the duration of the activity. Activities were classified as walking (structured walking for transport or in leisure time), moderate-intensity PA (MPA) or vigorous-intensity PA (VPA), based on their energy requirements (MPA: 3–6 MET; VPA: >6 MET; MET = metabolic energy equivalent task)⁽²⁶⁾.

Long-form IPAQ

The self-administered long-form IPAQ was used to assess PA at work, transport-related PA, domestic and gardening activities, and PA during leisure time in a usual week. Total times engaged in walking, MPA and VPA, all expressed in min/week, were computed according to the guidelines (www.ipaq.ki.se). The IPAQ has been shown to be a valid and reliable instrument for measuring PA in Europe⁽¹⁵⁾ and in Flanders, Belgium⁽²⁷⁾.

Short-form IPAQ

PA was assessed using the self-administered short-form IPAQ, which provides information on the time spent walking, in MPA and in VPA (min/week) in a usual week. This version of the IPAQ has been found to be valid and reliable⁽¹⁵⁾.

Baecke questionnaire

In the Baecke questionnaire⁽¹⁶⁾, responses were scored on a five-point scale and resulted in three different indices reflecting PA during work (work index), PA during leisure time excluding sport (LT index) and PA during sport activities (sport index). The summation of the three indices was defined as the overall PA index (total PA index). Good validity of the Baecke questionnaire for the assessment of PA was found in the past^(28,29).

Data analysis

Analyses were carried out using the SPSS for Windows statistical software package version 12.0 (SPSS Inc., Chicago, IL, USA). Average daily steps were calculated and values above 20 000 steps/d were recorded as 20 000 steps/d to limit unrealistically high average step counts⁽³⁰⁾. Because of the non-normal distributions in PA data, Spearman correlation coefficients (r_s) were calculated to assess the relationship between step counts and questionnaire-based PA. The same technique was used to assess the correlations between PA data derived from the different questionnaires. Correlations were interpreted as low (<0.30), moderate (0.30–0.50) or high (>0.50). Differences in questionnaire-based PA between the five step count groups based on the pedometer indices⁽⁵⁾ were evaluated with multivariate ANOVA. First, the skewed questionnaire outcomes were log-transformed to approximate normal distributions. Parametric analyses were performed on the log-transformed data and adjusted for age, gender and employment status. *F* values, *P* values and partial η^2 , as a measure of the effect size, are reported. For reasons of clarity and comparability, the means and standard deviations of the non-transformed data are used in Table 3. Finally, cross-tabulations of the number of participants (not) reaching the step count thresholds of 7500, 10 000 and 12 500 steps/d and/or the guideline of 30 min MVPA/d were performed, and the percentages of agreement between the different PA recommendations were calculated based on the cross-tabulations. A *P* value ≤ 0.05 was considered as statistically significant.

Results

The correlations between step counts and questionnaire-based PA are shown in Table 2. Step counts correlated moderately with total PA ($r_s = 0.37$) and MPA ($r_s = 0.31$), and lowly with walking ($r_s = 0.19$) and VPA ($r_s = 0.25$), reported in the long-form IPAQ. Low correlations were found between step counts and total PA ($r_s = 0.28$), walking ($r_s = 0.15$) and VPA ($r_s = 0.20$) reported in the short-form IPAQ, except for MPA ($r_s = 0.33$). A low correlation was found between step counts and walking ($r_s = 0.10$) and VPA ($r_s = 0.16$) reported in the MLTPAQ; total PA ($r_s = 0.32$) and MPA ($r_s = 0.32$) reported in the MLTPAQ correlated moderately with step counts. Finally, step counts correlated moderately with the total PA index ($r_s = 0.44$), LT index ($r_s = 0.44$) and sport index ($r_s = 0.31$) of the Baecke questionnaire, except for the work index ($r_s = 0.19$). Correlations between PA reported in the different questionnaires are also presented in Table 2.

Table 3 shows the mean (SD) amounts of questionnaire-based PA (min/week) for the different step count groups divided according to the pedometer indices of Tudor-Locke and Bassett⁽⁵⁾. Significant differences between the

Table 2 Correlation matrix: Spearman correlation coefficient (95% CI)

	Step counts			Short-form IPAQ			MLTPAQ			Baecke Q	
	Total PA†	Walking	MPAT	Total PA†	Walking	MPAT	Total PA†	Walking	MPAT	VPA†	Total PA index
Long-form IPAQ											
Total PA†	0.37 (0.27–0.46)			0.73 (0.67–0.78)							
Walking	0.19 (0.08–0.30)	0.68 (0.62–0.74)					0.36 (0.26–0.45)	0.24 (0.13–0.34)			0.30 (0.20–0.40)
MPAT	0.31 (0.21–0.41)		0.56 (0.78–0.63)					0.40 (0.30–0.49)			
Short-form IPAQ											
Total PA†	0.25 (0.14–0.35)			0.78 (0.73–0.82)			0.31 (0.21–0.41)	0.20 (0.10–0.31)			0.31 (0.21–0.41)
Walking	0.28 (0.17–0.38)	0.15 (0.04–0.26)						0.41 (0.31–0.50)			0.40 (0.30–0.49)
MPAT	0.33 (0.23–0.43)		0.33 (0.23–0.43)								
MLTPAQ											
Total PA†	0.20 (0.09–0.31)										
Walking	0.32 (0.22–0.42)	0.10 (–0.01–0.21)									
MPAT	0.32 (0.22–0.42)		0.16 (0.05–0.27)								
Baecke Q											
Total PA index	0.16 (0.05–0.27)										
Work index	0.44 (0.35–0.53)										
LT index	0.19 (0.08–0.30)										
Sport index	0.44 (0.35–0.53)										
	0.31 (0.20–0.41)										

IPAQ, International Physical Activity Questionnaire (short-form, short version of the questionnaire; long-form, long version of the questionnaire); MLTPAQ, Minnesota Leisure Time Physical Activity Questionnaire; PA, physical activity; MPAT, moderate physical activity; VPA, vigorous physical activity; LT, leisure time. †In min.

five step count groups were found for total PA, MPA and VPA reported in the long-form IPAQ, short-form IPAQ and MLTPAQ. For walking, only data based on the short-form IPAQ differed significantly between the step count groups. The work, leisure-time, sport and total PA indices of the Baecke questionnaire differed significantly between the five step count groups.

In total, 80.6% of participants reached 7500 steps/d, 45.0% reached 10 000 steps/d and 39.4% reached 12 500 steps/d, whereas 30 min MVPA/d was reached by 85.4% according to the long-form IPAQ, 84.8% according to the short-form IPAQ and 68.1% according to the MLTPAQ (Fig. 1).

Using the long-form IPAQ, the step count thresholds of 7500, 10 000 and 12 500 steps/d and the 30 min MVPA/d recommendation were in agreement in respectively 77.7%, 54.4% and 48.7% of the participants (see Table 4). Based on the short-form IPAQ, respectively 76.5%, 51.8% and 48.1% of participants were well placed. Using the MLTPAQ, the recommendations were in agreement in respectively 71.9%, 53.1% and 51.9% of the participants. Some 89.2% of the participants reaching 7500 steps/d, 94.2% of those reaching 10 000 steps/d and 93.4% of those reaching 12 500 steps/d reached 30 min MVPA/d, according to the long-form IPAQ. With the short-form IPAQ these figures were respectively 88.0%, 90.6% and 91.8%. Of the participants, 74.8% reaching 7500 steps/d, 73.4% reaching 10 000 steps/d and 75.4% reaching 12 500 steps/d reached 30 min MVPA/d based on the MLTPAQ.

Discussion

The first aim of the present study was to evaluate the associations between step counts and questionnaire-based PA. Objectively measured step counts correlated positively with the subjectively reported PA levels. Significant, positive associations between step counts and questionnaire-based PA were also found in previous studies^(9,31) conducted in the USA, where walking is one of the most commonly reported forms of PA⁽³²⁾. A median correlation of $r = 0.33$ (range: 0.02–0.94) was found between step counts and self-reported PA in a review of mostly non-European studies⁽³³⁾. Surprisingly, in the present study, a low correlation was found between step counts and walking reported in the long-form IPAQ, short-form IPAQ and MLTPAQ. An explanation for this remarkable finding could be the lack of sensitivity of questionnaires⁽³⁴⁾ to detect walking, resulting in under-reporting of walking behaviour. Bassett *et al.*⁽³⁵⁾ also found that subjects underestimated their daily walking distance in the College Alumnus Questionnaire, compared with pedometer-based values. Furthermore, different types of walking were assessed through the various questionnaires. The long-form IPAQ asked about the combination of walking in various domains (i.e. work, transport,

Table 3 Differences in questionnaire-based physical activity (PA) across step count groups: healthy adult volunteers, Flanders, Belgium

Step count range Participants	Pedometer index										F (P)	Post hoc analysest	η^2	
	Sedentary		Low active		Somewhat active		Active		Highly active					
	<5000 n 12	5000–7499 n 55	7500–9999 n 84	10 000–12 499 n 61	≥12 500 n 64	Mean	SD	Mean	SD	Mean				SD
Total PA (min/week)														
Long-form IPAQ	462	405	722	676	1069	820	1329	804	1542	904	12.4***	a, h	0.14	
Short-form IPAQ	243	268	838	927	772	733	1286	1080	1458	1113	7.4***	c, h	0.09	
MLTPAQ	199	194	347	309	416	279	524	430	485	326	8.2***	c, e, f, g, h	0.10	
Walking (min/week)														
Long-form IPAQ	81	52	247	361	317	392	450	459	432	485	1.8 ^{NS}		0.02	
Short-form IPAQ	58	63	558	702	424	538	754	753	634	702	5.4***	c, e, g, h	0.07	
MLTPAQ	103	116	76	73	106	116	110	131	108	121	1.2 ^{NS}		0.02	
MPA (min/week)														
Long-form IPAQ	303	331	334	371	525	441	611	434	733	476	7.8***	a, b, c, h	0.10	
Short-form IPAQ	82	116	147	294	210	290	349	539	545	620	6.8***	a, b, c, d, h	0.08	
MLTPAQ	104	107	255	298	292	238	373	371	336	253	7.4***	c, f, g, h	0.05	
VPA (min/week)														
Long-form IPAQ	55	178	129	194	155	258	215	302	263	352	4.6***	c, d, f, h	0.06	
Short-form IPAQ	47	144	113	188	135	243	172	272	206	257	2.7*	a, c, d, f, h	0.04	
MLTPAQ	3	10	17	31	27	67	37	65	46	80	4.2**	a, c, d, h	0.05	
Baecke questionnaire														
Work index	1.8	0.5	1.8	0.6	2.0	0.6	2.2	0.7	2.3	0.6	6.9***	a, b, f, h	0.09	
LT index	1.9	0.6	2.3	0.5	2.5	0.6	2.6	0.6	2.6	0.7	6.3***	f, g, h	0.09	
Sport index	1.4	1.2	1.9	1.3	1.9	1.2	2.2	1.2	2.5	2.1	3.2*	a, b, c, d, f, h	0.05	
Total PA index	5.1	1.9	6.0	1.5	6.4	1.4	7.0	1.5	7.4	1.5	10.4***	a, c, h	0.13	

IPAQ, International Physical Activity Questionnaire (long-form, long version of the questionnaire; short-form, short version of the questionnaire); MLTPAQ, Minnesota Leisure Time Physical Activity Questionnaire; MPA, moderate physical activity; VPA, vigorous physical activity.

Significance of differences between the five groups: * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$ (NS, $P > 0.05$).

†Some *post hoc* pairwise comparisons were not significantly different: a, sedentary–low active NS; b, sedentary–somewhat active NS; c, low active–somewhat active NS; d, low active–highly active NS; e, low active–highly active NS; f, somewhat active–active NS; g, somewhat active–highly active NS; h, active–highly active NS.




Fig. 1 Percentage of participants reaching the step count of 7500 steps/d, 10 000 steps/d and 12 500 steps/d, and the standard of 30 min of moderate to vigorous physical activity (MVPA) per day, as assessed by three different questionnaires: the long version and short version of the International Physical Activity Questionnaire (long-form IPAQ, short-form IPAQ) and the Minnesota Leisure Time Physical Activity Questionnaire (MLTPAQ)

home, leisure time) while the MLTPAQ assessed leisure/pleasure walking and walking to work only. The pedometer measured all ambulatory activity throughout the day.

Furthermore, differences in questionnaire-based PA between groups based on the pedometer indices of Tudor-Locke and Bassett⁽⁵⁾ were evaluated. Results

showed that the classification of activity levels based on step counts also highlighted differences in self-reported levels of PA (low to medium effect sizes). The more active participants were, based on step counts, the higher the levels of total, MPA and VPA reported in the long-form IPAQ, short-form IPAQ and MLTPAQ. The application of

Table 4 Number of participants reaching both, one or neither of two types of physical activity recommendations: 7500 steps/d, 10 000 steps/d, 12 500 steps/d and the 30 min of moderate to vigorous physical activity (MVPA) per day standard, as assessed by three different questionnaires

30 min MVPA/d based on...	7500 steps	
	Not reached	Reached
Long-form IPAQ		
Not reached	18	27
Reached	42	223
Short-form IPAQ		
Not reached	17	30
Reached	43	220
MLTPAQ		
Not reached	36	63
Reached	24	187
	10 000 steps	
	Not reached	Reached
Long-form IPAQ		
Not reached	37	8
Reached	133	131
Short-form IPAQ		
Not reached	34	13
Reached	136	126
MLTPAQ		
Not reached	62	37
Reached	108	102
	12 500 steps	
	Not reached	Reached
Long-form IPAQ		
Not reached	37	8
Reached	151	114
Short-form IPAQ		
Not reached	37	10
Reached	151	112
MLTPAQ		
Not reached	69	30
Reached	119	92

IPAQ, International Physical Activity Questionnaire (long-form, long version of the questionnaire; short-form, short version of the questionnaire); MLTPAQ, Minnesota Leisure Time Physical Activity Questionnaire.

these findings is that individuals who engaged more in MPA and VPA accumulated more steps per day. Again, unexpectedly, this was not the case for walking reported in the long-form IPAQ and MLTPAQ. Only walking in the short-form IPAQ differed significantly between step count groups. All indices (work, leisure-time, sport and total PA) based on the Baecke questionnaire also increased gradually when step counts augmented. Concluding, step counts are capable of discriminating between total, MPA and VPA reported in the different questionnaires.

A third objective of the present study was to compare currently used PA guidelines, to show differences in the percentages of participants reaching the step count thresholds. Of the participants, 80.6% reached 7500 steps/d, 45.0% reached 10 000 steps/d and 39.4% reached 12 500 steps/d. The guideline of 30 min MVPA/d, on the other hand, was reached by 85.4% according to

the long-form IPAQ, followed by 84.8% according to the short-form IPAQ and by 68.1% according to the MLTPAQ. There are various possible explanations for this discrepancy. It is known that over-reporting of PA may occur when using self-reported questionnaires, including the IPAQ⁽⁷⁾. The lower percentage according to the MLTPAQ could be explained by the 1-year time frame of the questionnaire, which could cause recall biases. Also the fact that the MLTPAQ was interviewer-administered may have had an influence: research assistants were trained to detect over-reporting of PA⁽⁶⁾.

The current study also explored whether reaching different step count thresholds was sufficient to reach 30 min MVPA/d. When participants reached 7500 steps/d, the MVPA recommendation was reached in 89.2% (by long-form IPAQ; 88.0% by short-form IPAQ). Of participants, 94.2% (by long-form IPAQ; 90.6% by short-form IPAQ) of those reaching 10 000 steps/d reached 30 min MVPA/d. Participants reaching 12 500 steps/d reached 30 min MVPA/d in 93.4% (by long-form IPAQ; 91.8% by short-form IPAQ) of cases. All figures based on the MLTPAQ were lower (respectively 74.8%, 73.4% and 75.4%). However, these results indicated that step count thresholds are stringent enough. Le Masurier *et al.*⁽¹⁹⁾ found that 91% of women reaching the 10 000 steps/d standard accumulated more than 30 min MVPA/d based on accelerometer data. In the present study, it was easier for participants reaching the step count thresholds to reach 30 min MVPA/d, rather than the other way around. No more than half of the participants (respectively 49.6%, 48.1% and 48.6% based on the long-form IPAQ, short-form IPAQ and MLTPAQ) reaching 30 min MVPA/d reached 10 000 steps/d, for example. Welk *et al.*⁽⁹⁾ reported that participants who performed more than 30 min of PA daily reached at least 10 000 steps/d some 73% of the time. Tudor-Locke *et al.*⁽³⁶⁾ found that approximately 3000 steps are expected in 30 min of moderate-intensity ambulatory activity.

The present average step count level (12 100 steps/d) is clearly higher than that of a representative sample of Belgian adults (9700 steps/d)⁽²³⁾. A possible explanation could be the fact that the present study population is a convenience sample of mostly working volunteers, implying cautious interpretations and limited generalizability of the results.

The present results reflect that caution is needed when assessing PA standards, especially when using different methods. It is remarkable that despite moderate correlations between step counts and questionnaire-based PA, the percentages reaching the standards differ notably. The subjective nature of questionnaires, possibly causing over-reporting, can be an explanation of the discrepancy. To avoid this problem, a gold standard, such as doubly labelled water or accelerometers, could be used to determine if the 30 min MVPA/d guideline is effectively reached.

In summary, there is a modest relationship between step counts and questionnaire-based PA in the present European sample. Objective pedometers not only provide a measurement of walking but also give an indication of total, MPA and VPA. Even though pedometers alone cannot discriminate between the intensity of activities nor reflect the amount of time spent in specific intensity PA categories, they provide sufficient information to be valuable in PA assessment in large, free-living populations. Less agreement was found between the currently used PA guidelines (30 min MVPA/d and step count thresholds), suggesting that caution is needed when comparing active quantities in different samples based on different methods.

Acknowledgements

The authors declare that they have no competing interests. The study was not funded.

K.A.D.C. participated in the data collection, analysis and interpretation, and led the writing of the paper. G.M.C. and I.M.D.B. participated in the data collection, conceptualization of the study and provided substantive feedback on the manuscript. All authors have read and approved the final manuscript.

The authors would like to acknowledge all subjects participating in the study.

References

- Blair SN, Cheng Y & Holder S (2001) Is physical activity or physical fitness more important in defining health benefits? *Med Sci Sports Exerc* **33**, S379–S399.
- Haskell W, Lee I, Pate R, Powell K, Blair S, Franklin B, Macera C, Heath G, Thompson P & Bauman A (2007) Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* **39**, 1423–1434.
- Hatano Y (1993) Use of the pedometer for promoting daily walking exercise. *Int Counc Health Phys Educ Recreation* **29**, 4–8.
- Hatano Y (1997) Prevalence and use of pedometer. *Res J Walking* **1**, 45–54.
- Tudor-Locke C & Bassett DR (2004) How many steps per day are enough? Preliminary pedometer indices for public health. *Sports Med* **34**, 1–8.
- Montoye HJ, Kemper HC, Saris WH & Washburn RA (1996) *Measuring Physical Activity and Energy Expenditure*. Champaign, IL: Human Kinetics.
- Sallis J & Saelens BE (2000) Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q Exerc Sport* **71**, 1–14.
- Bassett DR, Ainsworth BE, Legget SR, Mathien CA, Main JA, Hunter DC & Duncan GE (1996) Accuracy of five electronic pedometers for measuring distance walked. *Med Sci Sports Exerc* **28**, 1071–1077.
- Welk GJ, Differding JA, Thompson RW, Blair SN, Dziura J & Hart P (2000) The utility of the Digi-Walker step counter to assess physical activity patterns. *Med Sci Sports Exerc* **32**, S481–S488.
- Shephard FK (1989) Assessment of physical activity and energy needs. *Am J Clin Nutr* **50**, 1195–1200.
- Hornbuckle LM, Bassett DR & Thompson DL (2005) Pedometer-determined walking and body composition variables in African-American women. *Med Sci Sports Exerc* **37**, 1069–1074.
- Sequeira MM, Rickenbach M, Wietlisbach V, Tullen B & Schutz Y (1995) Physical activity assessment using a pedometer and its comparison with a questionnaire in a large population study. *Am J Epidemiol* **142**, 989–999.
- Bouchard C, Shepard RJ, Stephens T, Sutton JR & McPherson BD (1990) *Exercise, Fitness, and Health. A Consensus of Current Knowledge*. Champaign, IL: Human Kinetics.
- Taylor HL, Jacobs DR, Schucker B, Knudsen J, Leon AS & Debacker G (1978) Questionnaire for the assessment of leisure time physical activities. *J Chronic Dis* **31**, 741–755.
- Craig CL, Marshall AL, Sjöström M *et al.* (2003) International Physical Activity Questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* **35**, 1381–1395.
- Baecke JA, Burema J & Frijters JE (1982) A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* **36**, 936–942.
- Wilde BE, Sidman CL & Corbin CB (1002) A 10,000-step count as a physical activity target for sedentary women. *Res Q Exerc Sport* **72**, 411–414.
- Hultquist CN, Albright C & Thompson DL (2005) Comparison of walking recommendations in previously inactive women. *Med Sci Sports Exerc* **37**, 676–683.
- Le Masurier GC, Sidman CL & Corbin CB (2003) Accumulating 10,000 steps: does this meet current physical activity guidelines? *Res Q Exerc Sport* **74**, 389–394.
- Miller R & Brown W (2004) Meeting physical activity guidelines and average daily steps in a working population. *J Phys Act Health* **1**, 217–225.
- Crouter SE, Schneider PL, Karabulut M & Bassett DR (2003) Validity of 10 electronic pedometers for measuring steps, distance and energy cost. *Med Sci Sports Exerc* **35**, 1455–1460.
- Tudor-Locke C, Lind KA, Reis JP, Ainsworth BE & Macera CA (2003) A preliminary evaluation of a pedometer-assessed physical activity self-monitoring survey. *Field Methods* **15**, 1–17.
- De Cocker K, Cardon G & De Bourdeaudhuij I (2007) Pedometer-determined physical activity and its comparison with the International Physical Activity Questionnaire in a sample of Belgian adults. *Res Q Exerc Sport* **78**, 429–437.
- Miller R, Brown W & Tudor-Locke C (2006) But what about swimming and cycling? How to ‘count’ non-ambulatory activity when using pedometers to assess physical activity? *J Phys Act Health* **3**, 257–266.
- De Backer G, Kornitzer M, Sobolski J, Dramaix M, Degré S, de Marneffe M & Denolin H (1981) Physical activity and physical fitness levels of Belgian males aged 40–55 years. *Cardiology* **67**, 110–128.
- Ainsworth BE, Haskell WL, Whitt MC *et al.* (2000) Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* **32**, 498–504.
- Vandelanotte C, De Bourdeaudhuij I, Sallis J, Philippaerts R & Sjöström M (2005) Reliability and validity of a computerized and Dutch version of the International Physical Activity Questionnaire (IPAQ). *J Phys Act Health* **2**, 63–75.
- Pols MA, Buenodemesquita HB, Ocke MC, Wentick CA, Kemper HCG & Collette HJA (1995) Validity and repeatability of a modified Baecke questionnaire on physical activity. *Int J Epidemiol* **24**, 381–388.

29. Philippaerts FM, Westerdorp KR & Lefevre J (1999) Doubly labeled water validation of three physical activity questionnaires. *Int J Sports Med* **20**, 284–289.
30. Tudor-Locke C, Ham S, Macera C, Ainsworth BE, Kirtland KA, Reis JP & Kimsey Jr CD (2004) Descriptive epidemiology of pedometer-determined physical activity. *Med Sci Sports Exerc* **36**, 1567–1573.
31. Bassett DR, Schneider PL & Huntington GE (2004) Physical activity in an old order Amish community. *Med Sci Sports Exerc* **36**, 79–85.
32. Crespo CJ, Keteyian SJ, Heath GW & Sempos CT (1996) Leisure-time physical activity among US adults. *Arch Intern Med* **17**, 423–473.
33. Tudor-Locke C, Williams JE, Reis JR & Pluto D (2002) Utility of pedometers for assessing physical activity: convergent validity. *Sports Med* **32**, 795–808.
34. Ainsworth BE, Leon AS, Richardson MR, Jacobs DR & Paffenbarger RS (1993) Accuracy of the college alumnus physical activity questionnaire. *J Clin Epidemiol* **46**, 1403–1411.
35. Bassett DR, Cureton AL & Ainsworth BE (2000) Measurement of daily walking distance – questionnaire versus pedometer. *Med Sci Sports Exerc* **32**, 1018–1023.
36. Tudor-Locke C, Sisson SB, Collova T, Lee SM & Swan PD (2005) Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly health population. *Can J Appl Physiol* **30**, 666–676.